

*Original Scientific Article***EFFECT OF THE ADMINISTRATION OF PGF₂ α ANALOGUE TO EXTENDED BOAR SEMEN ON SPERM MOTILITY, MORPHOLOGY AND KINEMATIC PARAMETERS**

Mihail Chervenkov¹, Teodora Ivanova², Paulina Taushanova¹,
Rossen Stefanov¹, Boyko Georgiev¹

¹*Bulgarian Academy of Sciences, Institute of Biology and Immunology of Reproduction,
73 Tzarigradsko Shosse Bul., 1113 Sofia, Bulgaria*

²*Bulgarian Academy of Sciences, Institute of Biodiversity and Ecosystem Research,
23 Acad. G. Bonchev Str. 1113 Sofia, Bulgaria*

Received 23 December 2016; Received in revised form 24 February 2017; Accepted 20 March 2017

ABSTRACT

The addition of prostaglandin F₂ α (PGF₂ α) to boar semen prior to insemination improves the conception and farrowing rates in sows. It is accepted that this is due to increased myometrial contractility, which improves the spermatozoa movement. However, there are limited data about the effect of the exogenous PGF₂ α analogs on sperm motility parameters and morphology. The aim of the current study was to define if there are changes in motility, morphology and kinematic parameters of spermatozoa on 1st and 24th hour after addition of PGF₂ α analogue to extended boar semen. A total of 18 ejaculates, obtained from clinically healthy boars were diluted 1:3 in semen extender, and each of them was separate into four aliquots, 50 ml each. PGF₂ α was added to 3 of them in concentrations of 6, 12 and 25 μ g/ml, and the fourth served as untreated control. The motility, kinematic parameters and morphology of spermatozoa were evaluated on 1st and 24th hours after addition of PGF₂ α . There was no significant difference in sperm morphology, total and progressive motility between the untreated and treated groups. There was however a significant decrease in the rapid velocity and some of the kinematic parameters (VCL, VSL and VAP) in the group treated with 25 μ g/ml compared to the control at the 1st hour after PGF₂ α treatment, which (except for the rapid velocity) persisted to the 24th hour. The results indicate that addition of Oestrophan (Bioveta,CZ) to the extended boar semen did not improve the sperm motility, morphology and kinematic parameters of the spermatozoa.

Key words: boar semen, PGF₂ α , Oestrophan, motility, kinematic parameters

INTRODUCTION

Artificial insemination (AI) is very important for modern pig farming. In comparison to natural mating, AI allows obtaining a greater litter size per sow by using sires with better fertility and genetic qualities. Among the main goals of the specialists in pig reproduction is achieving an even better

conception rate after AI. A lot of research has been focused on the addition of different compounds to the semen extenders, which can protect and preserve sperm motility and integrity for a prolonged period, or can enhance the sperm movement, through the female reproductive tract. One of the compounds with such potential is prostaglandin F₂ α (PGF₂ α). In sows, PGF₂ α and its analogues have been widely used for synchronization of estrus (1), inducing of parturition (2) or abortion (3) and to reduce the weaning-to-estrus interval (4). The administration of PGF₂ α in boars can improve the training of sexually active boars for semen collection (5), and increase the volume and the total number of sperm in the sperm rich fraction (6). Another asset of PGF₂ α is the fact that it can increase myometrial contractility (7). Subsequently, this will facilitate the sperm movement through the female reproductive

Corresponding author: Dr. Mihail Chervenkov, PhD
E-mail address: vdmchervenkov@abv.bg
Present address: Bulgarian Academy of Sciences, Institute of Biology and Immunology of Reproduction, 73 Tzarigradsko Shosse Bul., 1113 Sofia, Bulgaria
Phone: +359 2 872 00 18 *Fax:* +359 2 872 00 22

Copyright: © 2017 Chervenkov M. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

Available Online First: 25 April 2017

Published on: 15 October 2017

<https://doi.org/10.1515/macvetrev-2017-0017>

tract. Kos and Bilkei (8) demonstrated that boar semen supplemented with PGF2 α immediately before insemination improves the conception and farrowing rates, as well as regular returns to estrus in the inseminated sows. The amount of endogenous PGF2 α in boar semen is less than 100 pg/ml, which is insufficient to increase myometrial contractility (9), so addition of exogenous PGF2 α is needed for obtaining this beneficial effect. For the reasons mentioned above, the supplementation of boar semen with PGF2 α or its analogues can be beneficial for the success of AI in pigs.

However, there is a question arising: What is the effect of PGF2 α or its analogues on sperm motility parameters, when added to extended boar semen? In an experiment with boars from the Landrace breed, the addition of PGF2 α led to an increase of sperm motility to over 80% while the control group had motility of 65-70% (10). In another study, the administration of PGF2 α in semen from Pietrain boar did not increase the semen motility parameters (11).

The aim of current study was to define if there are changes in motility, morphology and kinematic parameters of spermatozoa on 1st and 24th hour after addition of PGF2 α analogue (cloprostenol, Oestrophan, Bioveta, CZ) to extended boar semen.

MATERIAL AND METHODS

Semen collection

The ejaculates (n=18) were collected manually (gloved-hand technique) from clinically healthy Danube white breed boars, at the age of 2.5 years. The semen was collected twice per week in accordance with regulatory requirements for breeding in pigs.

Sperm motility and kinematic parameters

Sperm concentration, total motility, progressive motility, rapid velocity as well as kinematic parameters - Curvilinear Velocity (VCL); Straight-line Velocity (VSL); Average Path Velocity (VAP); Linearity (LIN); Straightness (STR); Wobble (WOB); Amplitude of lateral head displacement (ALH) and Beat/Cross Frequency (BCF) were evaluated by a computer-assisted sperm analyzer (CASA) (SCA, Microptic, Spain)

Sperm morphology

Sperm smears were prepared on preparation glass and stained by Sperm Blue (Microptic, Spain), performed according to the manufacturer's

protocol. The sperm morphology was evaluated under microscope (Nikon Eclipse E200, Japan) under a bright field with magnification of $\times 40$.

Experimental design

Each ejaculate was evaluated under a microscope immediately after obtaining and those not complying with standard requirements were discarded. Subsequently, they were diluted 1:3 in semen extender "Sredetz" and transferred to the laboratory. Each ejaculate was separated into four aliquots, 50 ml each. PGF2 α analogue (cloprostenol, Oestrophan, Bioveta, CZ) was added to 3 of them in concentrations of 6, 12 and 25 $\mu\text{g/ml}$, and the fourth served as untreated control. The samples were stored at 15°C before the evaluations. The motility, kinematic parameters and morphology of spermatozoa were assessed at 1st and 24th hours after addition of PGF2 α .

Statistical analysis

Values were presented as mean \pm standard deviation. The statistical analyses were performed by software R 2.10. All data was tested for variance using ANOVA and then experimental treatments were compared to the untreated control using Dunnett t-test ($p \leq 0.05$).

RESULTS

The present study was focused on the evaluation of the effect of PGF2 α analogue cloprostenol on motility characteristics and morphology of spermatozoa, when added to extended boar semen. The motility and morphology of the spermatozoa at the 1st and the 24th hour after PGF2 α analogue administration are presented in Table 1. There was a decrease in the percent of spermatozoa with progressive motility in the experimental groups compared to the control in both time points, which however was not statistically significant ($p > 0.05$). The rapid velocity also decreased in time more rapidly in PGF2 α treated groups than in control, but statistical significance was observed only at the 1st hour post treatment with 25 $\mu\text{g/ml}$ ($p = 0.019$). There was no significant difference in the percent of spermatozoa with normal morphology between the untreated and treated groups, at both time periods.

There was no statistically significant difference in the kinematic parameters of boar spermatozoa in the control and the groups treated with 6 or 12 $\mu\text{g/ml}$

Table 1. Effect of PGF2 α analogue on motility and morphology of boar spermatozoa at the 1st and 24th hour after administration in the extended semen (n=18)

Treatment		Total motility		Progressive motility		Rapid velocity		Normal morphology	
		Mean%	SD \pm	Mean%	SD \pm	Mean%	SD \pm	Mean%	SD \pm
Control	1 h	92,16	0,58	69,81	3,78	83,65	9,22	88,67	2,08
	24 h	91,63	1,52	58,44	2,33	90,68	3,55	87,00	1,73
6 μ g	1 h	91,10	1,85	67,19	6,62	60,84	22,49	88,00	1,73
	24 h	91,53	1,39	55,17	2,84	89,55	4,26	86,67	2,08
12 μ g	1 h	92,13	0,53	69,65	3,41	66,42	3,96	86,67	0,58
	24 h	90,33	2,04	56,08	3,20	82,63	3,57	85,00	2,00
25 μ g	1 h	89,76	3,280	62,51	11,99	47,36*	4,16	85,67	3,06
	24 h	88,27	2,94	52,99	9,22	79,22	10,40	84,00	2,65

¹Normal morphology - spermatozoa without morphological defects in head, midpiece and tail region and without cytoplasmic droplets. SD- standard deviation. *-p<0.05

PGF2 α analogue (Table 2). The addition of 25 μ g/ml PGF2 α analogue to boar semen led to a statistically significant decrease in some of the kinematic parameters of the spermatozoa - VCL (p=0.025), VSL (p=0,002) and VAP (p=0,008) at the 1st hour after treatment, while the other LIN, STR, WOB,

BCF and ALH remained unchanged (Table 2). Similar results were observed at the 24th hour, with some changes in the level of significance - VCL (p=0.025), VSL (p=0,02) and VAP (p=0,015). Again, the other kinematic parameters remained unchanged.

Table 2. Effect of PGF2 α analogue on kinematic parameters of boar spermatozoa at the 1st and 24th hour after administration in the extended semen (n=18)

Treatment		VCL, μ m/s		VSL, μ m/s		VAP, μ m/s		LIN,%		STR,%		WOB,%		BCF, Hz		ALH, μ m	
		Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm
Control	1 h	109,55	13,17	28,91	2,40	55,72	6,39	26,47	1,02	52,02	1,69	50,88	0,32	7,95	0,55	4,27	0,51
	24 h	106,57	10,28	28,20	2,19	53,83	5,31	26,53	1,00	52,53	1,88	50,50	0,26	7,50	0,70	4,67	0,40
6 μ g	1 h	92,06	21,27	24,64	2,25	46,10	8,21	27,38	3,94	54,04	4,97	50,49	2,71	8,64	1,44	3,78	0,84
	24 h	108,93	2,39	27,57	1,04	54,60	2,09	25,30	0,56	50,47	0,61	50,17	1,52	8,17	0,85	4,63	0,12
12 μ g	1 h	94,70	5,07	26,21	1,02	47,46	2,56	27,71	1,45	55,28	2,18	50,12	0,87	8,06	0,97	3,73	0,21
	24 h	88,77	4,37	22,87	2,06	43,57	1,90	25,87	3,49	52,40	2,71	49,23	4,14	6,70	0,26	4,33	0,15
25 μ g	1 h	74,40*	3,50	19,91**	2,45	37,26**	2,16	26,74	2,65	53,30	3,44	50,10	2,15	6,52	1,07	3,70	0,02
	24 h	82,20*	13,60	20,27*	4,51	39,37*	7,42	24,50	1,45	51,30	3,08	47,83	2,06	6,10	0,82	4,03	0,51

SD-standard deviation. *p<0.05, **p<0.01, ***p<0.001

DISCUSSION

Our results indicate that administration of PGF2 α analogue (Oestrophan, Bioveta, CZ) at doses of 6 and 12 $\mu\text{g}/\text{ml}$ in extended boar semen did not affect the total and progressive motility and the percent of rapid velocity spermatozoa of Danube white boars, at the 1st and 24th hour after treatment. Similarly to our results, Maes et al. (11) found no improvement in motility and kinematic parameters of the spermatozoa after addition of PGF2 α (Dinolytic; Pharmacia Animal Health; 5 mg PGF2 α /ml) in doses of 2.5, 5 or 10 mg PGF2 α to 100 ml diluted sperm from Pietrain boars. In another study, Pandur and Pacala (10) found that the addition of 1 ml PGF2 α (Dinolytic®; Pfizer; 5 mg PGF2 α /ml) to 100 ml semen from Landrace boars, right after dilution, lead to increase in sperm motility. The difference between our findings and those of Pandur and Pacala (10) might be contributed to the different breed and the different extender that were used, the different time of motility evaluation (immediate PGF2 α addition in theirs, and at the 1st and 24th hour after treatment in our experiment) or the fact that the sperm motility assessment in their work, was performed not by CASA but by a technician on a phase-contrast microscope.

Yeste et al. (12) supplemented extended semen from Pietrain boars with PGF2 α (Dinolytic®, Pharmacia, 5mg PGF2 α /ml) in concentrations of 0.625, 1.25, 2.50, 5, 10, 12.50, 25 and 50 mg PGF2 α to 100 ml semen, and assed the sperm quality immediately after treatment, and after 1, 3, 6 and 10 days. They found that treatments with PGF2 α in concentrations higher than 12.5 mg/100 ml were cytotoxic, while the others did not damage boar spermatozoa. None the less, the addition of PGF2 α at 5 mg/100 ml to sperm diluted in BTS extender may maintain sperm viability and motility better after 6 days of cooling, compared with control at the same time.

Also It was found that administration of 0.25 or 2.5 μg cloprostenol per 1ml diluted boar semen had no impact on boar sperm motility, while concentrations higher than 5 μg had a negative effect (13).

In our study the concentration of 25 $\mu\text{g}/\text{ml}$ PGF2 α analogue (Oestrophan, Bioveta, CZ) leads to a significant decrease in some of the kinematic parameters of the sperm at the 1st and the 24th hour after treatment. Additionally, the same concentration affects negatively the percent of rapid velocity spermatozoa at 1st hour after treatment. The other treatments have no negative effect on the sperm motility, morphology and kinematic parameters.

Our findings suggested that PGF2 α analogue cloprostenol supplementation to extended boar semen has no positive effect on sperm motility and kinematic parameter. The obtained data is in agreement with most of the other research work in the area (11, 12, 13, 14), despite the fact that we used cloprostenol (Oestrophan, Bioveta, CZ) – a synthetic analogue of PGF2 α , while in some of the other research work dinoprost tromethamine was used – a naturally occurring prostaglandin F2 α . The above mentioned suggests that even if the origin of the PGF2 α analogues is different, the effect on boar semen is similar. The decrease in kinematic parameters and subsequently in rapid velocity which occurs with higher concentration of cloprostenol may be due to different factors like mitochondrial transformation or induced sperm membrane damage. Similarly, the concentration of cloprostenol higher than 300 $\mu\text{g}/\text{ml}$ of diluted bull semen leads to decrease in sperm motility and even induces sperm membrane damage and permeability (14).

CONCLUSION

The results indicate that addition of PGF2 α analogue cloprostenol to the extended boar semen did not improve the sperm motility, morphology and kinematic parameters of the spermatozoa from Danube white boars. The positive effect of PGF2 α on the conception and farrowing rates, demonstrated by other research work which included supplementation of PGF2 α analogues to boar semen prior AI, can be contributed to other factors (e.g. increase myometrial motility), but not to improvement of sperm intrinsic movement.

CONFLICT OF INTEREST STATEMENT

The authors declared that they have no potential conflict of interest with respect to the authorship and/or publication of this article

ACKNOWLEDGEMENT

This research work was partially supported by Contract NoBG051PO001-3.3.06-0059, funded by the European Social Fund and the Operational Program Human Resources Development (2007-2013) and co-funded by the Ministry of Education and Science of the Republic of Bulgaria

REFERENCES

1. Noguchi, M., Kashiwai, S., Itoh S., Okumura H., Kure K., Suzuki C., Yoshioka K. (2013). Reproductive hormone profiles in sows on estrus synchronization using estradiol dipropionate and prostaglandin F(2 α)-analogue and the reproductive performance in female pigs on commercial farms. *J Vet Med Sci.* 75(3): 343–348.
<https://doi.org/10.1292/jvms.12-0022>
PMid:23131781
2. Einarsson, S., Fischler, M., Karlberg, K. (1981). Induction of parturition in sows using prostaglandin F2 α or the analogue cloprostenol. *Nordic Vet Med.* 33, 354–358.
PMid:6948275
3. Pressing, A.L., Dial, G.D., Stroud, C.M., Almond, G.W., Robison, O.W. (1987). Prostaglandin - induced abortion in swine: endocrine changes and influence on subsequent reproductive activity. *Am J Vet Res.* 48(1): 45-50.
PMid:3826841
4. Morrow, M., Britt, J., Belschner, A., Neeley, G., O'Carroll, J. (1996). Effect of injecting sows with prostaglandin F2 α immediately postpartum on subsequent reproductive performances. *Swine Health Prod.* 4, 73–78.
5. Estienne, M., Harper, A. (2000). PGF2 α facilitates the training of sexually active boars for semen collection. *Theriogenology.* 54, 1087–1092.
[https://doi.org/10.1016/S0093-691X\(00\)00417-9](https://doi.org/10.1016/S0093-691X(00)00417-9)
6. Hashizume, T., Niwa, T. (1984). Effect of administration of prostaglandin F2 α (PGF2 α) on the properties of sperm rich fraction of boar semen. *Japan. J. Anim. Reprod.* 30, 182-185.
<https://doi.org/10.1262/jrd1977.30.182>
7. Cheng, H., Althouse, G.C., Hsu, W.H. (2001). Prostaglandin F2 α added to extended boar semen at processing elicits in vitro myometrial contractility after 72 h storage. *Theriogenology* 55, 1901–1906.
[https://doi.org/10.1016/S0093-691X\(01\)00531-3](https://doi.org/10.1016/S0093-691X(01)00531-3)
8. Kos, M., Bilkei, G. (2004). Prostaglandin F2 α supplemented semen improves reproductive performance in artificially inseminated sows. *Animal Reproduction Science* 80, 113–120.
[https://doi.org/10.1016/S0378-4320\(03\)00135-0](https://doi.org/10.1016/S0378-4320(03)00135-0)
9. Cheng, H., Althouse, G.C., Hsu, W. H. (2003). Concentrations of endogenous prostaglandin F2 α in boar semen and effect of a 72-h incubation period on exogenous prostaglandin F2 α concentration in extended boar semen. *Prostaglandins & Other Lipid Mediators* 70, 285–290.
[https://doi.org/10.1016/S0090-6980\(02\)00140-5](https://doi.org/10.1016/S0090-6980(02)00140-5)
10. Pandur, I. D., Pacala, N. (2012). Sperm motility after the addition of prostaglandin F2 α to the Landrace boar diluted semen. *Scientific Papers Animal Science and Biotechnologies* 45 (1): 222-225.
11. Maes, D., Mateusen, B., Rijsselaere, T., De Vliegher, S., Van Soom, A., de Kruif, A. (2003). Motility characteristics of boar spermatozoa after addition of prostaglandin F2 α . *Theriogenology* 60, 1435–1443.
[https://doi.org/10.1016/S0093-691X\(03\)00132-8](https://doi.org/10.1016/S0093-691X(03)00132-8)
12. Yeste, M., Briz, M., Pinart, E., Sancho, S., Garcia-Gil, N., Badia, E., Bassols, J., Pruneda, A., Bussalleu, E., Casas, I., Bonet, S. (2008). Boar spermatozoa and prostaglandin F2 α quality of boar sperm after the addition of prostaglandin F2 α to the short-term extender over cooling time. *Animal Reproduction Science* 108, 180–195.
<https://doi.org/10.1016/j.anireprosci.2007.08.008>
PMid:17897798
13. Kozumplik J, Martinek J. (1986). The effect of Oestrophan Spofa (synthetic analog of prostaglandin F2 α) added to the insemination dose on pregnancy and fertility in sows. *Vet Med (Praha).* 31(4): 227–232.
14. Fayed A. (1996). Effect of prostaglandin F2 α and methylxantines on enzymic release of bull epididymal spermatozoa in vitro. *Contraception* 53(3): 181–184.
[https://doi.org/10.1016/0010-7824\(96\)00008-X](https://doi.org/10.1016/0010-7824(96)00008-X)

Please cite this article as: Chervenkov M., Ivanova T., Taushanova P., Stefanov R., Georgiev B. Effect of the administration of PGF2 α analogue to extended boar semen on sperm motility, morphology and kinematic parameters. *Mac Vet Rev* 2017; 40 (2): 125-129.
<https://doi.org/10.1515/macvetrev-2017-0017>