



Original Scientific Article

RABIES VIRUS NUCLEOPROTEIN: LARGE-SCALE EXPRESSION
IN PROKARYOTIC SYSTEM

Leandro Daniel Picotto¹, Carlos Javier Panei^{2,3}, Marcelo Ricardo Pecoraro²,
Guillermo Hernán Sguazza²

¹Reproductive Biotechnology, National Agricultural Technology Institute,
(INTA), Ruta 226 km 73.5, 7620, Balcarce, Argentina

²Virology Laboratory, Faculty of Veterinary Sciences,
National University of La Plata, 60&118, CC 296, 1900, La Plata, Argentina

³National Scientific and Technical Research Council (CONICET), Argentina

Received 5 April 2024; Received in revised form 13 August 2024; Accepted 15 August 2024

ABSTRACT

Rabies is controlled by mass animal vaccination campaigns. Cats, dogs, and wild animals (e.g., bats) are large reservoirs of this virus and can pose a significant threat to the human health, especially in the developing countries. The nucleoprotein of the rabies virus is of great scientific interest since it has the potential to generate immunity in animals and can be used as for immunochemical diagnostics. The study aimed to test a large-scale expression of the rabies N protein in a prokaryotic system. The recombinant N protein was successfully expressed and purified. It was immunologically recognized by specific antibodies and was able to induce the production of specific antibodies in a mouse immunization assay. These encouraging results indicate that the recombinant N protein can be evaluated as an antigen for the development of a subunit vaccine or for a diagnostic assay.

Key words: rabies virus, nucleoprotein, *Escherichia coli*, large-scale expression

INTRODUCTION

Rabies is one of the oldest viral diseases recorded in the world causing a fatal form of encephalomyelitis in humans and animals when no treatment is administered. The existence of large reservoirs of rabies viruses in stray dogs and a lot of wild animal species, such as bats, is a serious health problem for humans and domestic animals (1).

Rabies is currently controlled through mass animal vaccination campaigns and reduction of the stray animal population through a program for the rescue and responsible adoption of stray dogs

and cats (2). Vaccination is achieved by means of different commercial vaccines developed for human and veterinary medicine.

Vaccines can be produced in nervous tissue and cell culture (3). The production of biological subunit vaccines for rabies virus could be achieved under low biosafety level and could reduce the risk of infection for people who work with this highly-deadly virus (4).

The rabies virus belongs to the genus *Lyssavirus* within the family *Rhabdoviridae*. This virus has a negative-sense single-stranded ribonucleic acid (RNA) genome which encodes five proteins in the following order: nucleoprotein (N), phosphoprotein, matrix protein, glycoprotein (G), and RNA-dependent RNA polymerase (5, 6).

The N protein of the Pasteur strain rabies virus contains 450 amino acids and has approximate size of 51 kDa (7). It can generate immunity against rabies virus when administered to animals (8). This protein is the main target for T-helper lymphocytes (9) and has several characteristics that make it function as an exogenous superantigen

Corresponding author: Dr. Carlos Javier Panei, PhD

E-mail address: javierpanei@fcv.unlp.edu.ar

Present address: Virology Laboratory, Faculty of Veterinary Sciences,
National University of La Plata 60&118, CC 296, 1900, La Plata, Argentina
Phone: +542214824956

Copyright: © 2024 Picotto L.D. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

Available Online First: 3 September 2024

Published on: 15 October 2024

<https://doi.org/10.2478/macvetrev-2024-0026>

(10). The N protein is useful as viral antigen for immunochemical diagnostics of the rabies infection. Also, the antibodies produced by viral antigens have been used as diagnostic reagents to distinguish between rabies and rabies-related viruses (11). The expression and cloning in *E. coli* has a significant advantage in yields and purity (12). The study aimed to test a large-scale expression of the rabies virus N protein in bacterial system for the potential use as antigen and/or immunogen design.

MATERIAL AND METHODS

Strains

The CVS strain of the rabies virus, provided by the Instituto Biológico Dr. Tomás Perón of La Plata, Buenos Aires, Argentina, was used for the extraction of viral RNA and for western blot assays. The pQE-30 UA vector (Qiagen, Germany) was used for cloning and expression of N protein in *E. coli* strain M15 (Qiagen, Germany). This vector presents a His-tag at the N-terminus to facilitate the purification of the proteins of interest. The Luria Bertani (LB) medium with 100 µg/ml of ampicillin and 25 µg/ml of kanamycin was used for the screening of *E. coli* colonies transformed with the pQE-30 vector containing the N gene (pQE-30-N) and for the expression of the N protein in *E. coli* (with Isopropyl β-D-1-thiogalactopyranoside, IPTG, 1 mM).

Primer design and amplification of the N gene by RT-PCR

To amplify a segment of the N gene, a set of primers were designed from a multiple alignment with several sequences of the rabies virus obtained by BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). In the design, the cleavage sites with the restriction enzymes *Bam*HI and *Hind*III were added in forward (NF: GTCTCGAGGATCCATGGATGCCGACAA) and reverse (NR: CGGAAGCTTATGAGTCACTCGAATATG) primers, respectively. After RNA extraction of the CVS strain of rabies virus, the reverse transcription (RT) reaction was carried out with the reverse transcriptase enzyme of the Moloney Murine Leukemia virus (MMLV) (Promega, USA). The PCR reaction was carried out using cDNA, the NF and NR designed primers, and the high-fidelity enzyme Platinum SuperFi DNA Polymerase (Invitrogen, USA). The reaction was carried out under the following thermal cycling conditions: 94 °C for 3 min of initial denaturation; 94 °C for 45 s, 60 °C for 45 s, and 72 °C for 1.5 min (30 cycles); and a final extension of 10 min at 72 °C. In order

to confirm the amplification of the N gene, an electrophoresis on a 1.5% agarose gel with 10 µl of the PCR product was carried out.

Ligation of the N gene in the pQE-30 vector and E. coli transformation

The N gene and pQE-30 vector were digested with the restriction enzymes *Bam*HI and *Hind*III (Promega, USA). The ligation was carried out following the protocol described for the ExpressLink™ T4 DNA Ligase kit (Invitrogen, USA). This ligation mixture was used to transform electrocompetent bacteria (*E. coli* M15) by electroporation. The transformation mix was plated on LB-Amp-Kan plates and incubated overnight at 37 °C. To confirm the presence of the N gene, 12 colonies were randomly selected and checked by colony-PCR with the NF and NR primers. The colony-PCR products were analyzed by 1.5% agarose gel electrophoresis. Finally, to verify the open reading frame of the cloned fragment, the plasmid DNA of one of the colony-PCR positive clones was purified and sequenced, using the sequencing primers included in the pQE-30 kit.

Expression of the N protein

With the colony-PCR positive clones, cultures 1/100 diluted in 5 ml of LB-Amp-Kan medium were performed and incubated at 37 °C overnight, with stirring at 100 rpm until an optical density (OD)_{600nm} between 0.5 and 0.7. The cultures were induced with 1 mM IPTG for 2 h at 37 °C. Total proteins were extracted from each culture and the expression level of N protein was analyzed by sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS-PAGE) and Western Blot (WB). A negative control from culture of M15 bacteria containing the empty pQE-30 vector was used. To optimize the concentration of IPTG and the induction time to generate the greatest amount of N protein, bacterial cultures were performed in 20 ml of LB-Amp-Kan with 0.5 and 1 mM of IPTG. From the beginning of the cultures, 2 ml were taken every hour (up to 3 h) from each flask. To compare the expression level of N protein of each culture, the total proteins of pellet samples were analyzed by SDS-PAGE and WB.

Rabbit anti-rabies polyclonal antibodies

For N protein expression level assays by WB, we produced a rabbit anti-rabies polyclonal antibody by using the veterinary rabies vaccine provided by the Pasteur Institute of Buenos Aires, Argentina. The immunization protocol was carried out in a rabbit,

which was immunized by subcutaneous injections with 1 ml of vaccine on days 0, 7, and 14. At 21 days post-inoculation, the rabbit was bled to obtain serum. As secondary antibody was used an anti-rabbit IgG (Sigma A9044, USA), conjugated with horseradish peroxidase.

Scale-up and purification

A batch culture was carried out in a 7.5-l BioFlo 310 Benchtop Bioreactor fermenter (New Brunswick-Scientific, Edison, NJ, USA), maintaining the flask conditions with the highest level of N protein expression. To perform the seed culture, 150 ml of fresh LB-Amp-Kan medium was inoculated with 1 ml of an overnight culture of the selected clone. After overnight incubation at 37 °C with shaking, the culture was used to inoculate 3.85 l of new fresh LB-Amp-Kan medium in the fermenter (initial OD_{600nm} ~ 0.1). The temperature and pH were fixed at 37 °C and 7 °C respectively. The agitation was programmed to achieve aerobic growth of the microorganism and guarantee an oxygen concentration of at least 30% of the saturation value during all the process. Once the culture reached an OD_{600nm} of 0.6, it was induced with IPTG at a final concentration of 1 mM and kept for 3 h in culture. Culture samples were taken every hour, and, after 3 h, the culture was centrifuged at 3,000 x g for 20 min at 4 °C, discarding the supernatant. Finally, the pellet was analyzed by SDS-PAGE and WB. The purification was carried out by immobilized metal affinity chromatography in a 10 ml column loaded with a matrix of nitrilotriacetic acid and nickel (NTA-Ni) (Bio-Rad, USA), following the manufacturer's recommendations. The eluted fraction containing the purified protein was dialyzed against phosphate-buffered saline, sterilized by filtration, and its concentration was estimated by means of an SDS-PAGE with a bovine serum albumin (BSA) standard and, using the TotalLab program.

Evaluation of the recombinant N protein

To evaluate whether the recombinant N protein induce the production of specific antibodies against the same native protein of rabies virus, mice were immunized by the intramuscular (IM), subcutaneous (SC) and intraperitoneal routes (IP). After inoculation, serum samples were obtained from each animal and analyzed by WB. A single formulation containing 10% aluminum hydroxide as adjuvant (equivalent to 10 mg/ml aluminum content) and 10 µg/ml of the recombinant N protein was used. The veterinary rabies vaccine, produced by the Pasteur Institute of Buenos Aires

in nervous tissue, was used as antigenicity control by the intraperitoneal (IP) route. A duplicate immunization protocol was carried out at 0 and 14 days by the intramuscular (IM), subcutaneous (SC), and IP routes. The amount of protein used in each route was determined according to previous reports (13). At 28 days after the first inoculation, the mice were bled by cardiac puncture.

A WB was carried out with the CVS strain as antigen to analyze the presence of antibodies against the native N protein in serum samples. After the blotting and blocking steps, the polyvinylidene difluoride membrane was cut into thin strips and used individually with 1/50 dilutions of the serum from each mouse. After incubation for 90 min and three washes in washing solution, the membrane was incubated for 90 min with a 1/1,000 dilution of an anti-mouse IgG serum conjugated with horseradish peroxidase produced in goats (Sigma cat. A9040). Finally, after additional three washes, the presence of bound antibodies was detected using a solution containing 3,3'-diaminobenzidine (DAB) (Merk, Germany) and hydrogen peroxide.

Ethical committee evaluation

The experimental procedures developed in this study comply with international regulations (Institute of Laboratory Animal Resources, 1996) and were approved by the Institutional Committee for Care and Use of Laboratory Animals (CICUAL) of the Faculty of Veterinary Sciences of UNLP (Exp.600-004104/08-001. Res. 129/09).

RESULTS

RT-PCR amplification

The amplification of the N gene was successful because an approximate size of 1.4 kb was observed (Fig. 1A), which coincided with the size for the segment expected (1373 bp). Over 100 colonies were obtained after transforming *E. coli* M15 cells with the pQE-30-N vector (N gene ligated to pQE-30 vector). Screening of the transformation of 12 colonies was carried out, showing a positive result in 6 colonies (1, 2, 3, 6, 9, and 12), confirming the pQE-30-N construction (Fig. 1B). After sequencing, the start codon of the N gene was checked in frame with a His-tag in pQE-30-N. Also, analyzing the sequence by means of a local alignment by BLAST, more than 99 % sequence identity was obtained with the N gene of the viral strains Pasteur (GU992320.1), SAD-1 (EF206718.1), SAD-BERN (LN713619.1), and ERA (AB781935.1).




Figure 1. A- RT-PCR of the *N* gene. Lane 1: Thermo Scientific™ GeneRuler 1 kb Plus DNA Ladder; lane 2: RT-PCR product. B- Screening of *E. coli* colonies with pQE-30-N. Lane 1: Thermo Scientific™ GeneRuler 1 kb Plus DNA Ladder; lanes 2 to 13: pQE30-N candidate colonies (colonies 1 to 12); lane 14: positive amplification control




Figure 2. A- Western blot of small-scale cultures. Lane 1: Pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); lane 2: Inactivated rabies virus; lane 3: negative control I (*E. coli* with empty pQE-30); lane 4: negative control II (*E. coli* without plasmid); lanes 5 to 10: colonies 1 to 6 with pQE-30-His-N. A 1:100 dilution of Rabbit anti-rabies polyclonal antibody produced in our laboratory and 1:1.000 dilution of anti-rabbit IgG (Sigma A9044, USA) were used. B- Culture optimization, SDS-PAGE. Lane 1: Pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); lane 2: negative control (*E. coli* with empty pQE-30), 3 h induction with 1 mM IPTG. Lanes 3 to 8: clone 6 pQE-30-N-His in different conditions; lane 3: 3 h induction with 0.5 mM IPTG; lane 4: 3 h induction with 1 mM IPTG; lane 5: 1 h induction with 0.5 mM IPTG; lane 6: 1 h induction with 1 mM IPTG; lane 7: 2 h induction with 0.5 mM IPTG; lane 8: 3 h induction with 1 mM IPTG. C- Culture optimization, western blot. Lane 1: inactivated CVS strain of rabies virus; lane 2: pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); lane 3: negative control (*E. coli* with empty pQE-30), 3 h induction with 1 mM IPTG. Lanes 4 to 9: clone 6 pQE-30-N-His in different conditions; lane 4: 1 hour induction with 0.5 mM IPTG; lane 5: 1 h induction with 1 mM IPTG; lane 6: 2 h induction with 0.5 mM IPTG; lane 7: 2 h induction with 1 mM IPTG; lane 8: 3 h induction with 0.5 mM IPTG; lane 9: 3 h induction with 1 mM IPTG. A 1:100 dilution of Rabbit anti-rabies polyclonal antibody produced in our laboratory and 1:1.000 dilution of anti-rabbit IgG (Sigma A9044, USA) were used

Expression of recombinant N protein

The protein expression level assay by Western blot (WB) was conducted on the culture supernatant of the six colony-PCR positive clones (clones 1 to 6), as shown in Fig. 2A. Among these clones, colony n° 6 exhibited recognition of His-N by specific antibodies against the rabies virus. Therefore, this clone was selected for further optimization of IPTG concentration and induction time required for scale-up. Subsequent analysis using SDS-PAGE (Fig. 2B) demonstrated that the expression of His-N was highest when induced with 1 mM IPTG for a duration of 3 h. WB analysis (Fig. 2C) was performed to confirm these results, and it revealed visible expression after a 2-hour induction period. Moreover, it was observed that higher levels of expression were achieved using 1 mM IPTG instead of 0.5 mM. Comparing the amount of His-N produced at both a 2-hour and a 3-hour induction period with the use of 1 mM IPTG showed greater

expression at the latter time point. Based on these findings, it was determined that optimal conditions for scale-up involve using an IPTG concentration of 1 mM and inducing for a duration of 3 h. The purity and expression level of His-N in the scaled culture were qualitatively compared, observing them before and after purification (Fig. 3A). Furthermore, the concentration of N-His of the purified fraction was estimated at 30 mg/l (Fig. 3B).

Immunological assays

This qualitative test confirmed the presence of specific antibodies against native rabies N protein in the serum of the animals inoculated by the three routes with the recombinant N protein. The level of specific antibodies against native rabies N protein produced in the serum of the animals inoculated by the SC and IP routes was similar than that observed in animals immunized with the reference rabies vaccine used in Argentina (Fig. 4).




Figure 3. A- Western blot of the culture in a fermenter and purification fractions. Lane 1: *E. coli* with pQE-30-His-N, 3 h of induction with 0.5 mM IPTG; lane 2: negative control (*E. coli* with empty pQE-30 plasmid), 3 h of induction with 1 mM IPTG; lane 3: washing fraction of purification (pH=6.3); lane 4: inactivated CVS strain of rabies virus; lane 5: Pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); lane 6: elution fraction of purification (pH=5.9); lane 7: elution fraction of purification (pH=4.5). A 1:100 dilution of Rabbit anti-rabies polyclonal antibody produced in our laboratory and 1:1.000 dilution of anti-rabbit IgG (Sigma A9044, USA) were used. B- SDS-PAGE of purified His-N. Lane 1: elution fraction of purification (pH=4.5); lane 2: Pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); lanes 3 to 7: 10, 20, 40, 60 and 80 mg/l of BSA



Figure 4. Western blot of the mouse sera inoculated with the recombinant N protein. Strip 1: Pre-stained protein molecular weight marker (PageRuler™ Thermo Scientific™, USA); strip 2: control serum 1 IP immunized with the reference vaccine; strip 3: mouse serum 1 IM immunized; strip 4: mouse serum 2 IM immunized; strip 5: mouse serum 1 SC immunized; strip 6: mouse serum 2 SC immunized; strip 7: mouse serum 1 IP immunized; strip 8: mouse serum 2 IP immunized; strip 9: control serum 2 IP immunized with the reference vaccine

DISCUSSION

Rabies is one of the oldest recorded diseases affecting humanity, dating back more than 2,000 years. Sporadic outbreaks of rabies in companion animals are increasingly frequent around the world (14). This increases the risk of transmission to domestic animals, and consequently to humans, since they tend to interact with bats. The option to control the virus in affected areas is by a mass vaccination plan for animals that can act as viral vehicles and a reduction of the stray animal population. Adverse reactions may include neurological complications which are attributed to the myelinated tissue in the vaccine produced in suckling mice (15), hence they have been discouraged by the WHO. Considering that rabies is a disease that mainly affects developing countries, an efficient and inexpensive solution is desirable. Recombinant or synthetic soluble antigens have been found to be safe and low-cost (16).

This rabies virus genome encodes five proteins in the following order: N, P, M, G, and L (5, 6). The most immunogenic and antigenic are N and G (17, 18). The G-protein ectodomain stimulates neutralizing antibody production to protect vaccinated animals from rabies virus infection by inducing T lymphocytes and T helper cells (9, 19, 20, 21). It has been extensively expressed in various heterologous systems, including yeast (22). The N protein from different rhabdoviruses showed highly conserved characteristic with shared group-specific antigenic determinants (23). Therefore, these proteins are great candidates to

be used as antigens in diagnostic aims (24) and/or as immunogen in triggering cellular immune responses against rabies (10).

The aim of this work was to test large-scale expression of the rabies virus N protein in bacterial system with implications to be potentially used as a design for antigen and/or immunogen. The obtained recombinant *E. coli* clones containing the N gene were subjected to a small-scale expression assay. Culture clones with the highest level of expression were selected. The IPTG concentration and the induction time to produce the highest level of N protein in flasks were optimized. The scale-up was carried out in a fermenter with the same conditions to improve the amount of N protein produced.

Plasmid stability is influenced by several factors such as copy number, the size of inserted DNA, and culture conditions, including temperature and growth rate (25, 26). Higher growth rates are associated with increased plasmid stability, while cell density can impact the instability of expression plasmids during induction. Therefore, to address this issue, our strategy involved conducting IPTG induction at low cell density (in exponential phase) to prevent the loss of expression plasmids in a short time (27).

The recombinant N protein produced and purified by affinity chromatography was specifically recognized by antibodies specific against the native N protein of the rabies virus. The recombinant N protein stimulated the production of specific antibodies against the native viral N protein in a similar level as the one reached the reference vaccine.

CONCLUSION

The recombinant N protein from rabies virus was produced in *E. coli* and was purified by affinity chromatography. It was immunologically recognized and was able to produce specific antibodies against native rabies N protein in immunological assay. In combination with the recombinant glycoprotein, it can be used for development of subunit vaccine or a diagnostic kit.

CONFLICT OF INTEREST

The authors declare that they have no known conflict of interest in the conduction of the current study.

ACKNOWLEDGMENTS

This study was funded by National University of La Plata, Argentina “(Programa de Incentivos a Docentes Investigadores, Proyecto 11/V224)”.

AUTHORS' CONTRIBUTION

LDP performed conceptualization, writing original draft, review and editing, conducted the investigation, methodology and visualization. CJP made the conceptualization, review and editing, visualization and formal analysis. MRP performed the review and editing. GHS conducted the investigation, review and editing of the paper.

REFERENCES

- Martinez, L. (2000). Global infectious disease surveillance. *Int J Infect Dis.* 4(4): 222-228.
[https://doi.org/10.1016/S1201-9712\(00\)90114-0](https://doi.org/10.1016/S1201-9712(00)90114-0)
PMid:11231187
- Kumar, A., Bhatt, S., Kumar, A., Rana, T. (2023). Canine rabies: an epidemiological significance, pathogenesis, diagnosis, prevention, and public health issues. *Comp Immunol Microbiol Infect Dis.* 97, 101992.
<https://doi.org/10.1016/j.cimid.2023.101992>
PMid:37229956
- Natesan, K., Isloor, S., Vinayagamurthy, B., Ramakrishnaiah, S., Doddamane, R., Fooks, A.R. (2023). Developments in rabies vaccines: the path traversed from Pasteur to the modern era of immunization. *Vaccines (Basel)* 11(4): 756.
<https://doi.org/10.3390/vaccines11040756>
PMid:37112668 PMCID:PMC10147034
- Zhou, X., Wang, H., Zhang, J., Guan, Y., Zhang, Y. (2024). Single-injection subunit vaccine for rabies prevention using lentinan as adjuvant. *Int J Biol Macromol.* 254 (Pt 3): 128118.
<https://doi.org/10.1016/j.ijbiomac.2023.128118>
PMid:37977452
- Coslett, G.D., Holloway, B.P., Obijeski, J.F. (1980). The structural proteins of rabies virus and evidence for their synthesis from separate monocistronic RNA species. *J Gen Virol.* 49(1): 161-180.
<https://doi.org/10.1099/0022-1317-49-1-161>
PMid:7420062
- Conzelmann, K.K., Cox, J.H., Schneider, L.G., Thiel, H.J. (1990). Molecular cloning and complete nucleotide sequence of the attenuated rabies virus SAD B19. *Virology* 175(2): 485-499.
[https://doi.org/10.1016/0042-6822\(90\)90433-R](https://doi.org/10.1016/0042-6822(90)90433-R)
PMid:2139267
- Rahmahani, J., Suwarno, S., Yuniarti, W.M., Rantam, F.A. (2019). Antigenic site of nucleoprotein gene from Indonesian rabies virus isolates. *Vet World.* 12(5): 724-728.
<https://doi.org/10.14202/vetworld.2019.724-728>
PMid:31327911 PMCID:PMC6584851
- Dietzschold, B., Lafon, M., Wang, H., Otvos, L., Jr, Celis, E., Wunner, W. H., Koprowski, H. (1987). Localization and immunological characterization of antigenic domains of the rabies virus internal N and NS proteins. *Virus Res.* 8(2): 103 -125.
[https://doi.org/10.1016/0168-1702\(87\)90023-2](https://doi.org/10.1016/0168-1702(87)90023-2)
PMid:2445121
- Celis, E., Karr, R., Dietzschold, B., Wunner, W., Koprowski, H. (1988). Genetic restriction and fine specificity of human T cell clones reactive with rabies virus. *J Immunol.* 141(8): 2721-2728.
<https://doi.org/10.4049/jimmunol.141.8.2721>
PMid:2459225
- Lafon, M., Lafage, M., Martinez-Arends, A., Ramirez, R., Vuillier, F., Charron, D., Lotteau, V., Scott-Algara, D. (1992). Evidence for a viral superantigen in humans. *Nature* 358, 507-510.
<https://doi.org/10.1038/358507a0>
PMid:1386410
- Goto, H., Minamoto, N., Ito, H., Luo, T.R., Sugiyama, M., Kinjo, T., Kawai, A. (1995). Expression of the nucleoprotein of rabies virus in *Escherichia coli* and mapping of antigenic sites. *Arch Virol.* 140(6): 1061-1074.
<https://doi.org/10.1007/BF01315415>
PMid:7611878

12. Reddy, G.B., Singh, R., Singh, R.P., Singh, K.P., Gupta, P.K., Mahadevan, A., Shankar, S.K., et al. (2011). Molecular characterization of Indian rabies virus isolates by partial sequencing of nucleoprotein (N) and phosphoprotein (P) genes. *Virus genes* 43(1): 13-17.
<https://doi.org/10.1007/s11262-011-0601-0>
PMid:21452061
13. Harlow, E., Lane, D. (1988). *Antibodies: a laboratory manual*. NY: Cold Spring Harbor Laboratory Press
14. Gan, H., Hou, X., Wang, Y., Xu, G., Huang, Z., Zhang, T., Lin, R., et al. (2023). Global burden of rabies in 204 countries and territories, from 1990 to 2019: results from the Global Burden of Disease Study 2019. *Int J Infect Dis.*126, 136-144.
<https://doi.org/10.1016/j.ijid.2022.10.046>
PMid:36343866
15. Plotkin, S., Orenstein, W., Offit, P., Kathryn, M.E. (2018). *Plotkin's Vaccines - 7th Ed.* Elsevier
16. Minke, J.M., Audonnet, J.C., Fischer, L. (2004). Equine viral vaccines: the past, present and future. *Vet Res.* 35(4): 425-443.
<https://doi.org/10.1051/vetres:2004019>
PMid:15236675
17. Lyu, X. (2023). Rabies proteins functions and future directions. *Theoret Nat Sci.* 21, 53-58.
<https://doi.org/10.54254/2753-8818/21/20230812>
18. Hampson, K., Coudeville, L., Lembo, T., Sambo, M., Kieffer, A., Attlan, M., Barrat, J., et al. (2015). Estimating the global burden of endemic canine rabies. *PLoS Negl Trop Dis.* 9(4): e0003709.
<https://doi.org/10.1371/journal.pntd.0003709>
PMid:25881058 PMCID:PMC4400070
19. Morales, M.M., Rico, R.G., Gómez, O.J., Aguilar, S.A. (2006). Immunologic importance of the N protein in the rabies virus infection. *Rev Vet Méx.* 37(3): 351-367.
20. Loza-Rubio, E., & Rojas-Anaya, E. (2014). Edible Rabies vaccines. In: J. Howard, E. Hood (Eds.), *Commercial plant-produced recombinant protein products* (pp. 153-177). *Biotechnology in Agriculture and Forestry*, Vol. 68. Berlin, Heidelberg: Springer
https://doi.org/10.1007/978-3-662-43836-7_9
PMCID:PMC7120656
21. Johnson, N., Cunningham, A.F., Fooks, A.R. (2010). The immune response to rabies virus infection and vaccination. *Vaccine* 28(23): 3896-3901.
<https://doi.org/10.1016/j.vaccine.2010.03.039>
PMid:20368119
22. Picotto, L.D., Sguazza, G.H., Tizzano, M.A., Galosi, C.M., Cavalitto, S.F., Pecoraro, M.R. (2017). An effective and simplified DO-stat control strategy for production of rabies glycoprotein in *Pichia pastoris*. *Protein Expr Purif.* 132, 124-130.
<https://doi.org/10.1016/j.pep.2017.02.004>
PMid:28189633
23. Tursunov, K., Begaliyeva, A., Ingirbay, B., Mukanov, K., Ramanculov, E., Shustov, A., Mukantayev, K. (2017). Cloning and expression of fragment of the rabies virus nucleoprotein gene in *Escherichia coli* and evaluation of antigenicity of the expression product. *Iran J Vet Res.* 18(1): 36-42.
24. Qin, S., Volokhov, D., Rodionova, E., Wirblich, C., Schnell, M.J., Chizhikov, V., Dabrazhynetskaya, A. (2019). A new recombinant rabies virus expressing a green fluorescent protein: A novel and fast approach to quantify virus neutralizing antibodies. *Biol: J Int Assoc Biol Stand.* 59, 56-61.
<https://doi.org/10.1016/j.biologicals.2019.03.002>
PMid:30898479
25. Valentinotti, S., Srinivasan, B., Holmberg, U., Bonvin, D., Cannizzaro, C., Rhiel, M., Von Stockar, U. (2003). Optimal operation of fed-batch fermentations via adaptive control of overflow metabolite, *Control Eng Pract.* 11(6): 665-674.
[https://doi.org/10.1016/S0967-0661\(02\)00172-7](https://doi.org/10.1016/S0967-0661(02)00172-7)
26. Babaeipour, V., Shojaosadati, S.A., Robotjazi, S.M., Khalilzadeh, R., Maghsoudi, N. (2007). Over-production of human interferon- γ by HCDC of recombinant *Escherichia coli*. *Process Biochem.* 42(1): 112-117.
<https://doi.org/10.1016/j.procbio.2006.07.009>
27. Babaeipour, V., Mofid, M.R., Khanchezar, R., Faraji, F., Abolghasemi, S. (2017). Bench-scale Overproduction and Purification of recombinant GCSF in *Escherichia coli* fed-batch process. *J Appl Pharm Sci.* 7(8): 149-155.