

*Original Scientific Article***PET OWNERS' KNOWLEDGE, ATTITUDES, AND PRACTICES REGARDING
ANTIMICROBIAL USE AND ANTIMICROBIAL RESISTANCE IN
COMPANION ANIMALS IN NORTH MACEDONIA**

Ivana Shikoska, Jane Vlahov, Boris Dimitrievski, Bojana Chapkunovska,
Ljubica Rashikj, Mirko Prodanov, Martin Nikolovski,
Elena Atanaskova Petrov, Aleksandar Cvetkovikj, Iskra Cvetkovikj

*Faculty of Veterinary Medicine-Skopje, Ss. Cyril and Methodius University in Skopje,
Lazar Pop Trajkov 5-7, 1000 Skopje, North Macedonia*

Received 12 March 2026; Received in revised form 28 April 2026; Accepted 23 June 2026

ABSTRACT

Antimicrobial use (AMU) in companion animals is influenced not only by veterinary prescribing practices but also by pet owners, whose decisions on medication administration, compliance, and expectations can impact antimicrobial stewardship and antimicrobial resistance (AMR) at the human-animal interface. This study aimed to assess pet owners' knowledge, attitudes, and practices regarding AMU and AMR in companion animals in North Macedonia. A cross-sectional, web-based questionnaire was completed by 410 pet owners in North Macedonia. Data was analysed using descriptive and inferential statistics. Of the respondents, 96.3% (395/410) acknowledged AMR as a serious health challenge, indicating high general awareness. However, this awareness did not consistently translate into accurate knowledge and recommended practices. For example, while 92.4% (379/410) identified bacteria as a target of antibiotic therapy, only 67.1% (275/410) chose bacteria as the only correct answer. Regarding interspecies transmission, 66.6% (273/410) agreed that companion animals may transmit antimicrobial-resistant bacteria to humans, while 60.2% (247/410) were previously aware of such transmission. Trust in veterinarians was high, but 36.1% (148/410) of respondents still expected antibiotic therapy for a wide range of clinical conditions. Additionally, 44.1% (181/410) reported administering antibiotics intended for human use to their pets, and 62.9% (258/410) reported keeping antibiotics at home "just in case." Knowledge of bacteriological testing was associated with adherence to the prescribed treatment duration. These findings indicate that, despite high general awareness of AMR, important gaps remain in pet owners' specific knowledge and practices. Pet owners should be considered an important target for antimicrobial stewardship efforts in small animal practice in North Macedonia.

Keywords: antibiotics, antimicrobial stewardship, awareness, small animal practice, AST

INTRODUCTION

Antimicrobial use (AMU) is a recognised driver of antimicrobial resistance (AMR), and human behaviour plays a central role in prescribing, dispensing, and consumption practices across medical disciplines, including veterinary medicine (1, 2). While the majority of veterinary antimicrobials are

used in food-producing animals, companion animals such as dogs and cats are increasingly recognised as potential reservoirs of antimicrobial-resistant bacteria due to their close contact with humans and the potential for interspecies transmission (3, 4).

A major concern in small animal practice is the frequent use of antimicrobials outside approved label directions, known as extra-label or off-label use (5). This practice is considerably more common in companion animals than in food-producing species. The main reasons are the limited number of antimicrobial products specifically authorised for pets and the presence of outdated or non-specific dosing guidelines (6). In the United States, extra-label antimicrobial use is legally allowed under veterinary supervision. In contrast, the European Union enforces stricter regulations, especially for critically

Corresponding author: Asst. Ivana Shikoska, DVM
E-mail address: ivana@fvm.ukim.edu.mk
Present address: Faculty of Veterinary Medicine-Skopje, Ss. Cyril and Methodius University in Skopje,
Lazar Pop Trajkov 5-7, 1000 Skopje, North Macedonia
Phone: +38978440195

Copyright: © 2026 Shikoska I. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

Available Online First: 2 October 2026

<https://doi.org/10.2478/macvetrev-2026-0026>

important antimicrobials (CIAs) for human health (7). Although the overall volume of antimicrobial use in companion animals is lower than in livestock, the use of critically important antimicrobials (CIAs) remains an important stewardship concern in small-animal practice (8, 9). In North Macedonia, our previous survey among small-animal veterinarians identified frequent prescribing of CIAs alongside limited implementation of bacteriological and antimicrobial susceptibility testing (10). However, the selective pressure exerted by the use of CIAs and human-approved antimicrobials in pets remains a concern (11). This is important because evidence suggests that companion animals may carry resistant bacteria of major public health importance. These include methicillin-resistant staphylococci, ESBL/AmpC-producing *Enterobacteriales*, and carbapenem-resistant Gram-negative bacteria, with potential exchange of resistant genes between humans and pets within the same household (12).

In small animal practice, AMU is shaped by two key human actors: veterinarians, who are responsible for clinical decision-making and prescribing, and pet owners, who administer treatments, influence demand, and set expectations about antimicrobial therapy (13). Although veterinarians remain authorised prescribers, their prescribing is shaped by veterinarian-, patient-, treatment-, and context-related factors (14). Both groups play key roles in AMU patterns in companion animals and are central to strategies that mitigate and control AMR (15). Recent studies show that pet owners apply different antibiotic standards for themselves and their pets, often with less knowledge and poorer practices in the veterinary context (16). Veterinarians also deal with clinical uncertainty, client pressure, and time constraints. These factors can lead to suboptimal prescribing, even when unnecessary (17). Together, these dynamics stress the need to address both prescriber behaviour and owner understanding within the One Health framework.

Our recent survey among small-animal veterinarians in North Macedonia identified concerning findings regarding AMU, including frequent owner pressure to prescribe antibiotics and owner-administered antibiotics prior to veterinary consultation (10). These findings suggest that pet owners may play an important role in shaping AMU in companion animals. However, data on pet owners' knowledge, attitudes, and practices regarding AMU and AMR in North Macedonia are currently lacking. Addressing this gap is important, as pet owners may influence treatment expectations, adherence to prescribed therapy, and antibiotic use at the household level. Therefore, this study aimed

to investigate pet owners' knowledge, attitudes, and practices regarding AMU and AMR in order to inform antimicrobial stewardship efforts in companion animals in North Macedonia.

MATERIAL AND METHODS

Study design

A cross-sectional, web-based survey was conducted among pet owners in North Macedonia from 25 August to 4 December 2025 to assess their knowledge, attitudes, and practices regarding AMU and AMR in companion animals.

Questionnaire development

A structured questionnaire was developed specifically for this study. It included items on demographic characteristics, general understanding of antibiotics and AMR, trust in veterinarians, expectations regarding antimicrobial therapy, adherence to prescribed treatment, storage and use of antibiotics at home, administration of antibiotics intended for human use, and acceptance or refusal of bacteriological testing. The questionnaire consisted of 25 items, presented as single-choice questions, multiple-choice questions, Likert-type scales, and optional free-text responses. It was pilot-tested on five pet owners to assess clarity. An English version of the complete questionnaire, including all questions and response options, is provided as Supplementary Material (<https://data.mendeley.com/datasets/vprxjwfzy9/1>).

Participants and recruitment

The survey was administered via Google Forms and could be completed on a mobile device or computer, with responses automatically stored. Eligible participants included adults (18 years or older) who owned at least one pet. Recruitment was conducted online through social media platforms (Facebook and Instagram), online pet-owner communities, and the official social media pages of veterinary clinics that voluntarily shared the survey link. Additional recruitment was conducted at the University Veterinary Hospital of the Faculty of Veterinary Medicine – Skopje, where visitors were invited to participate by scanning a QR code displayed in the reception area. Participation was voluntary and anonymous. The survey settings limited responses to one submission per IP address to reduce duplicate entries. Informed consent was obtained from all participants prior to the initiation of the survey. No personal or sensitive data were collected.

Ethical statement

This questionnaire-based study did not involve any intervention, handling, or experimental procedures involving animals. Participation of pet owners was voluntary and anonymous, and informed consent was obtained before completion of the questionnaire. The collected data were used exclusively for scientific research purposes, and no personal or sensitive data were collected.

Statistical analysis

Data were entered and managed in Microsoft Excel and analysed using R statistical software (version 4.4.3; R Foundation for Statistical Computing, Vienna, Austria). Categorical variables are presented as absolute frequencies and percentages. Trust in veterinarians, measured on a five-point ordinal scale, is additionally summarised as median and interquartile range (IQR). Associations between categorical variables were assessed using Pearson's chi-square test of independence or Fisher's exact test, as appropriate. Associations between the ordinal trust variable and binary outcomes were examined using the Mann-Whitney U test. All tests were two-tailed, and p-values <0.05 were considered statistically significant. Given the exploratory nature of the study, no correction for multiple comparisons was applied. Three multivariable binary logistic regression models were fitted for the following outcomes: (1) belief that companion animals may represent a source of antibiotic-resistant bacteria transmissible between

animals and humans; (2) adherence to prescribed antibiotic duration (dichotomised as "always" versus "not always"); and (3) agreement that antibiotics may be used without veterinary consultation. Candidate predictors were selected based on clinical relevance and bivariate screening and were entered using a forced-entry approach. Results are presented as adjusted odds ratios (aORs) with 95% confidence intervals (CIs). Age was initially categorized into six groups (18–24, 25–34, 35–44, 45–54, 55–64, and >64 years), as presented in Table 1. To explore potential age-related differences across selected outcomes, these categories were subsequently combined into two broader groups (18–34 years and ≥35 years) and compared using the same bivariate approach. Model calibration was assessed using the Hosmer–Lemeshow goodness-of-fit test.

RESULTS

Sociodemographic data of the respondents

A total of 410 pet owners were included in the analysis. The sample was predominantly female (79.8%). The most common age group was 25–34 years (41.0%), followed by 35–44 years (22.9%). Approximately one quarter of respondents (24.9%) reported a medical or veterinary educational background. Most participants (72.0%) owned a single pet type, while 28.0% reported owning multiple pet types. Detailed sociodemographic characteristics are presented in Table 1.

Table 1. Sociodemographic characteristics of the respondents (n = 410)

Variable	Category	Frequency (n)	Percentage (%)
Gender	Male	83	20.2
	Female	327	79.8
Age (years)	18–24	52	12.7
	25–34	168	41.0
	35–44	94	22.9
	45–54	65	15.9
	55–64	28	6.8
	>64	3	0.7
Medical background	Yes	102	24.9
	No	308	75.1
Pet ownership	Single pet type	295	72.0
	Multiple pet types	115	28.0

*Knowledge**Knowledge of antibiotics and antimicrobial resistance*

Most respondents (92.4%) selected bacteria as targets of antibiotic therapy; however, because multiple responses were permitted, 27.4% also selected viruses and/or fungi. Overall, 67.1% correctly identified bacteria only as the sole target of antibiotic therapy, as shown in Table 2.

Almost all respondents (96.3%) recognized AMR as a serious health challenge affecting humans, animals and the environment. Regarding

transmission awareness, 66.6% agreed that companion animals may represent a potential source of antibiotic-resistant bacteria transmissible between animals and humans. Prior awareness of interspecies transmission was reported by 60.2% of participants, whereas 39.8% indicated they were not aware of this possibility before completing the questionnaire. Knowledge of bacteriological testing/AST varied: 54.9% reported familiarity, 20.5% reported no knowledge and 24.6% were unsure. Knowledge-related findings are summarized in Table 2.

Table 2. Knowledge of antibiotics and antimicrobial resistance (n=410)

	Question	Category	Frequency	Percentage (%)
Antibiotic effectiveness knowledge	*Antibiotics are effective against infections caused by:	Bacteria	379	92.4
		Viruses	68	16.6
		Fungi	73	17.8
		Do not know	15	3.7
	Correct response	Bacteria only	275	67.1
Recognition of antimicrobial resistance	Antimicrobial resistance represents a serious health challenge for humans, animals, and the environment	Yes	395	96.3
		No	15	3.7
Transmission awareness (One Health)	Companion animals represent a potential source of antibiotic-resistant bacteria transmissible between animals and humans	Yes	273	66.6
		No	137	33.4
	Prior to completing this questionnaire, were you aware that antibiotic-resistant bacteria can be transmitted between animals and humans and vice versa?	Yes	247	60.2
		No	163	39.8
Diagnostic knowledge	Do you know what bacteriological testing/AST is?	Yes	225	54.9
		No	84	20.5
		Not sure	101	24.6

*multiple choice allowed

*Attitudes**Trust and communication with veterinarians*

Trust in veterinarians was generally high, with 54.4% assigning the maximum trust score (5) and 26.1% selecting a score of 4. The median trust score was 5 (IQR 4–5). A score of 3 was reported by 15.9%, while 3.7% selected scores 1–2.

Most respondents reported that their veterinarian explains the reasons for prescribing antibiotics (86.8%) and specifies which antibiotic is prescribed (82.7%), whereas 13.2% and 17.3%, respectively, indicated that such explanations were not provided (Table 3).

Table 3. Trust and communication with veterinarians (n=410)

Variable	Category	Frequency (n)	Percentage (%)
Trust in veterinarian (1–5 scale)	1	6	1.5
	2	9	2.2
	3	65	15.9
	4	107	26.1
	5	223	54.4
Veterinarian explains reasons for prescribing antibiotics	Yes	356	86.8
	No	54	13.2
Veterinarian specifies which antibiotic is prescribed	Yes	339	82.7
	No	71	17.3

Disagreement between owners and veterinarians and perceived responsibility for antibiotic prescribing

Most respondents (83.9%) reported no prior disagreement with their veterinarian regarding antibiotic therapy. Disagreement related to perceived unnecessary antibiotic prescription was reported by 12.4%, whereas 3.7% reported disagreement due to perceived failure to prescribe antibiotics when expected. Among respondents who experienced disagreement, 37.1% sought a second opinion, 42.9% did not, and 20.0% considered seeking one but did not pursue it.

The majority of respondents (88.3%) indicated that primary responsibility for appropriate antibiotic prescribing in companion animals lies with the veterinarian; 11.2% believed responsibility should be shared between the veterinarian and the pet owner, and 0.5% attributed primary responsibility to the pet owner alone.

Perceived contributors to AMR reflected a multifactorial understanding: 50.2% selected “all

listed factors.” Among individual contributors, overuse of antibiotics was identified by 42.4%, inappropriate use by 32.9%, and shortening/ prolonging treatment duration by 11.0%.

Practices

Antibiotic use behaviours and acceptance of diagnostic testing

Most respondents reported always adhering to the prescribed duration of antibiotic therapy (88.5%), while 11.5% reported occasional or complete non-adherence. Most respondents (93.9%) disagreed that antibiotics may be used without veterinary consultation; however, 44.1% reported using antibiotics intended for human use in their pets. In addition, 62.9% reported keeping antibiotics at home “just in case”, either always (22.2%) or sometimes (40.7%). Willingness to accept bacteriological testing/AST if recommended by a veterinarian was high (96.6%), with 3.4% reporting refusal, as shown in Table 4.

Table 4. Antibiotic use practices (n=410)

Variable	Category	Frequency (n)	Percentage (%)
Agreement that antibiotics may be used without veterinary consultation	Agree	25	6.1
	Disagree	385	93.9
Adherence to prescribed antibiotic duration	Always	363	88.5
	Sometimes	43	10.5
	Never	4	1.0
Use of human antibiotics in pets	Yes	181	44.1
	No	229	55.9
Keeping antibiotics at home “just in case”	Always	91	22.2
	Sometimes	167	40.7
	Never	152	37.1
Willingness to accept bacteriological testing if recommended	Yes	396	96.6
	No	14	3.4

Potential reasons for not accepting bacteriological testing/AST

The optional question on potential reasons for not accepting bacteriological testing/AST allowed multiple responses. As the question was conditional on hypothetical non-acceptance of testing, respondents who did not answer this item were included in the “none of the above” category.

Overall, 55.4% reported no potential reason for non-acceptance. Among the reported reasons, urgency of therapy was most frequent (15.6%), followed by financial reasons (11.5%) and fear of worsening clinical condition (10.2%). Less frequently reported reasons included long waiting time (4.6%) and considering testing unnecessary (2.7%) (Table 5).

Table 5. Potential reasons for not accepting bacteriological testing/AST (n=410)

Reason	n	%
Urgency of therapy	64	15.6
Financial reasons	47	11.5
Fear of worsening condition	42	10.2
Long waiting time	19	4.6
Testing unnecessary	11	2.7
None of the above (including non-response)	227	55.4

Multiple responses allowed

Owner expectations regarding antibiotic prescription

Respondents most frequently expected antibiotic prescription for urinary tract infections (56.1%) and respiratory infections (52.9%). Expectations were

also commonly reported for ear infections (45.6%) and eye infections (43.4%). Notably, 36.1% of respondents selected “all of the above,” indicating broad expectations for antibiotic use across clinical conditions. Because multiple responses were permitted, the reported categories are not mutually exclusive. Responses are shown in Fig 1.




Figure 1. Clinical conditions for which owners expect antibiotic prescription (n=410)

Factors associated with KAP-related outcomes

Determinants of correct knowledge of antibiotic effectiveness

Medical/veterinary background was significantly associated with correct knowledge of antibiotic effectiveness (“bacteria only”) ($\chi^2(1)=14.35$, $p<0.001$; Cramér’s $V=0.19$), although the effect size was small. Respondents with a

medical/veterinary background more frequently selected the correct response compared with those without such background (82.4% vs 62.0%) (Table 6). Recognition of AMR as a serious health challenge was also slightly more frequent among respondents with a medical/veterinary background (100% vs 95.1%; Fisher’s exact test, $p=0.028$; Cramér’s $V=0.11$), representing a small effect size (Table 6).

Table 6. Association between medical/veterinary background and correct knowledge of antibiotic effectiveness (n=410)

Medical background	Correct (Bacteria only) n (%)	Incorrect n (%)	Total
Yes (n=102)	84 (82.4%)	18 (17.6%)	102
No (n=308)	191 (62.0%)	117 (38.0%)	308
Total	275 (67.1%)	135 (32.9%)	410

Determinants of transmission beliefs

Prior awareness of interspecies transmission was strongly associated with the belief that companion animals may represent a potential source of antibiotic-resistant bacteria transmissible between animals and humans ($\chi^2(1)=80.87$, $p<0.001$; Cramér’s $V=0.44$). Notably, 40.5% of respondents without prior awareness still agreed

with the transmission concept. In multivariable binary logistic regression, prior awareness remained independently associated with transmission belief (aOR=7.83, 95% CI: 4.86–2.62, $p<0.001$), whereas AST knowledge was not associated after adjustment ($p=0.633$) (Table 7). The model demonstrated adequate calibration according to the Hosmer–Lemeshow goodness-of-fit test ($p=0.185$).

Table 7. Association between prior awareness and belief in animal-to-human transmission of antibiotic-resistant bacteria (n=410)

Prior awareness	Agree n (%)	Disagree n (%)	Total
Yes (n=247)	207 (83.8%)	40 (16.2%)	247
No (n=163)	66 (40.5%)	97 (59.5%)	163
Total	273 (66.6%)	137 (33.4%)	410

Determinants of adherence to prescribed antibiotic duration

Adherence was dichotomized as “always” versus “not always”. Knowledge of bacteriological testing/AST was significantly associated with adherence ($\chi^2(1)=19.83$, $p<0.001$; Cramér’s $V=0.22$),

representing a small-to-medium effect size; and respondents familiar with bacteriological testing/AST were more likely to report always adhering to the prescribed duration than those without such knowledge (95.1% vs 80.5%) (Table 8).

Table 8. Association between knowledge of bacteriological testing/AST and adherence to prescribed antibiotic duration (n=410)

Bacteriological testing/AST knowledge	Always n (%)	Not always n (%)	Total
Yes (n=225)	214 (95.1%)	11 (4.9%)	225
No/Unsure (n=185)	149 (80.5%)	36 (19.5%)	185
Total	363 (88.5%)	47 (11.5%)	410

Prior awareness of interspecies transmission was also associated with adherence ($\chi^2(1)=4.66$, $p=0.031$; Cramér’s $V=0.11$), though with a small effect size (Table 9). Trust in veterinarians did not differ between always-adherent and not-always-adherent respondents (median [IQR] 5 [4–5] vs 4 [4–5]; Mann–Whitney $U=9244$, $p=0.302$).

In multivariable binary logistic regression including knowledge of bacteriological testing/AST,

prior awareness of interspecies transmission, belief in animal-to-human transmission, age group, medical/veterinary background and trust in veterinarians, knowledge of bacteriological testing/AST remained independently associated with adherence (aOR=3.69, 95% CI: 1.64–8.33, $p=0.002$). The model demonstrated adequate calibration according to the Hosmer–Lemeshow goodness-of-fit test ($p=0.373$).

Table 9. Association between transmission awareness and adherence (n = 410)

Transmission awareness	Always n (%)	Not always n (%)	Total
Yes (n=247)	226 (91.5%)	21 (8.5%)	247
No (n=163)	137 (84.0%)	26 (16.0%)	163
Total	363 (88.5%)	47 (11.5%)	410

Determinants of permissive attitudes toward unsupervised antibiotic use

Agreement that antibiotics may be used without veterinary consultation was associated with keeping antibiotics at home (Fisher’s exact test,

$p=0.002$; Cramér’s $V=0.17$). Respondents who reported always keeping antibiotics at home were more likely to agree with unsupervised use (12.1%) compared with those who reported sometimes (1.2%) or never (7.9%) storing antibiotics (Table 10).

In multivariable binary logistic regression including storage of antibiotics and use of human antibiotics, keeping antibiotics at home remained independently associated with permissive attitudes toward antibiotic use without veterinary

consultation (aOR=6.39, 95% CI: 1.46–27.88, $p=0.013$). The model demonstrated adequate calibration according to the Hosmer–Lemeshow goodness-of-fit test ($\chi^2=0.15$, $p=0.700$).

Table 10. Association between storage of antibiotics and agreement that antibiotics may be used without veterinary consultation (n=410)

Storage of antibiotics	Agree n (%)	Disagree n (%)	Total
Always (n=91)	11 (12.1%)	80 (87.9%)	91
Sometimes (n=167)	2 (1.2%)	165 (98.8%)	167
Never (n=152)	12 (7.9%)	140 (92.1%)	152
Total	25 (6.1%)	385 (93.9%)	410

Age group differences

To explore potential age-related differences, respondents aged 18–34 years were compared with those aged ≥ 35 years. Younger respondents were less likely to report knowledge of bacteriological testing/AST (48.6% vs 62.1%; $\chi^2(1)=6.94$, $p=0.008$; Cramér's $V=0.13$) and less likely to report having used antibiotics intended for human use in their pets (39.1% vs 50.0%; $\chi^2(1)=4.49$, $p=0.034$; Cramér's $V=0.10$). No significant age-group differences were observed for adherence to prescribed antibiotic duration ($p=0.691$), storage of antibiotics at home ($p=0.847$), belief that companion animals may represent a source of antibiotic-resistant bacteria transmissible between animals and humans ($p=0.141$), prior awareness of interspecies transmission ($p=0.423$), or agreement that antibiotics may be used without veterinary consultation ($p=0.388$).

DISCUSSION

This cross-sectional survey provides new evidence on pet owners' knowledge, attitudes, and practices regarding AMU and AMR in companion animals in North Macedonia. Overall, respondents demonstrated high general awareness of AMR, with 96.3% recognizing it as a serious health challenge affecting humans, animals, and the environment. However, this general awareness did not consistently translate into stewardship-consistent knowledge and practices (18). Although 92.4% identified bacteria as the target of antibiotic therapy, only 67.1% selected bacteria as the sole

correct answer, indicating that misconceptions regarding antibiotic activity against viruses and fungi remain common. Similarly, while 66.6% agreed that companion animals may serve as a source of antibiotic-resistant bacteria transmissible between animals and humans, prior awareness of interspecies transmission was reported by only 60.2% of respondents. Together, these findings suggest that broad recognition of AMR does not necessarily imply a detailed understanding of appropriate antibiotic use or transmission-related risks.

Owner expectations and requests for antibiotic therapy have been identified as factors that may influence antimicrobial prescribing in veterinary practice (18). In the present study, trust in veterinarians was generally high, and most respondents reported that their veterinarian explained the reasons for prescribing antibiotics (86.8%) and specified which antibiotic was prescribed (82.7%). In addition, most respondents considered the veterinarian to be primarily responsible for appropriate antibiotic prescribing in companion animals. These findings indicate that veterinarians remain the central and most trusted source of treatment-related information in small animal practice, consistent with previous research showing that strong veterinarian-client relationships, clear communication, and shared decision-making are key drivers of acceptance of AMU recommendations (19). Data from Germany likewise showed high owner ratings of veterinarian communication on zoonoses and AMR, despite a gap between the information provided and the level of guidance desired by owners (20).

However, trust in veterinarians does not necessarily preclude expectations for antibiotic therapy. Scarborough et al. (21) reported that pet owners generally trusted veterinarians to act in their animals' best interests, even when no medication is prescribed, whereas Redding and Cole (22) found that many owners preferred their pet to receive antibiotics when the expected benefit of treatment was uncertain. This distinction is also reflected in our findings. Although respondents reported high trust and generally good communication with veterinarians, owner-reported expectations were not always aligned with antimicrobial stewardship principles. Specifically, 36.1% of respondents selected "all of the above" when asked about the clinical conditions for which they would expect an antibiotic prescription, suggesting that antibiotics may be perceived as appropriate across a broad range of clinical presentations, including conditions where they are not routinely indicated. Such expectations may contribute to pressure during consultations and to empirical antibiotic prescribing in small animal practice. This interpretation is consistent with findings from our previous national survey among small animal veterinarians in North Macedonia (10), in which 45.6% reported occasional and 35.1% frequent owner requests for antibiotic therapy, while owner-administered antibiotic use prior to veterinary consultation was also commonly reported (>90.0% overall; 54.4% occasional, 38.6% frequent).

Household-level access to antibiotics and owner-mediated AMU appear to be important drivers of antimicrobial exposure outside veterinary oversight (23). In the present study, although 93.9% of respondents disagreed with using antibiotics without veterinary consultation, 44.1% reported administering antibiotics intended for human use to their pets, and 62.9% reported keeping antibiotics at home "just in case." Similar findings have been reported in a UK-based survey of cat owners, where 17.3% of respondents reported keeping leftover antibiotics for future use, supporting the view that household availability of antimicrobials may facilitate unsupervised use (24). Together with findings from our previous survey among small-animal veterinarians in North Macedonia (10), these results suggest that owner-related influences on AMU may occur both during the prescribing encounter and after antibiotics become available in the household.

Keeping antimicrobials at home may facilitate treatment decisions outside veterinary oversight and promote more permissive attitudes toward

unsupervised antibiotic use (25). Similar patterns have been reported in a survey of Italian dog owners (23), where 21% reported administering oral antibiotics without veterinary consultation, and among these respondents, 65% used leftovers from previous prescriptions. In our study, household antibiotic storage was significantly associated with agreement that antibiotics may be used without veterinary consultation ($\chi^2=12.1$, $p=0.002$; Cramér's $V=0.17$), although the effect size was small. This association remained significant in multivariable analysis (aOR=6.4, 95% CI: 1.5–27.9, $p=0.013$), suggesting that household access to antimicrobials may be associated with greater acceptance of unsupervised antibiotic use.

The duration of antimicrobial therapy is an important component of appropriate AMU and antimicrobial stewardship (26). In the present study, 11.5% of respondents reported occasional or complete non-adherence to the prescribed therapy duration. Similar behaviour has been reported in other pet-owner populations; in one study, 92.4% of respondents indicated that they would discontinue treatment once clinical improvement was observed (27). This suggests that visible improvement may strongly influence owner decisions regarding treatment duration and highlights the need for stewardship-focused communication at the time of prescribing (28).

Notably, knowledge of bacteriological testing/AST was associated with better adherence to the prescribed duration of therapy in our study and remained an independent predictor in multivariable analysis. Respondents who were familiar with bacteriological testing/AST were more likely to always complete the prescribed antibiotic course than those who were not. Although the effect size of this association was modest (Cramér's $V=0.22$), the finding was consistent across bivariate and multivariable analyses, and may carry practical relevance in the context of stewardship. Although willingness to accept bacteriological testing when recommended by a veterinarian was high (96.6%), perceived urgency of therapy, financial constraints, and fear of clinical deterioration were identified as potential reasons for refusal. In clinical practice, such concerns may discourage diagnostic testing and favour empirical antimicrobial use in the absence of microbiological confirmation (29). When clinically appropriate, bacteriological examination and AST can guide treatment adjustment, including de-escalation from empirical broad-spectrum therapy to a narrower-spectrum antimicrobial (26).

In addition to owner-reported AMU practices, recent microbiological investigations conducted in companion animals in North Macedonia have demonstrated a substantial prevalence of AMR among both commensal and clinical bacterial isolates. Multidrug resistance was identified in 50.0% of *Escherichia coli* isolates obtained from shelter dogs, with 15.2% confirmed as ESBL/AmpC producers (30). Similarly, clinical *Staphylococcus* isolates from dogs showed multidrug resistance in 70.5% and methicillin resistance detected in 28.8% based on the presence of the *mecA* gene (31). In this context, owner awareness of transmission risk becomes particularly important. In the present study, 60.2% of respondents reported prior awareness of interspecies transmission, and this was associated with the belief that companion animals may represent a source of antibiotic-resistant bacteria transmissible between animals and humans, with a moderate effect size (Cramér's $V=0.44$). This proportion appears higher than that reported by Stein et al. (32), who found that only 21% of pet owners considered antimicrobial use in pets to pose a risk to humans. Together, these findings suggest that owner awareness and owner-mediated AMU practices are relevant not only to antimicrobial stewardship in companion animals, but also to the One Health risk of circulation and possible transmission of resistant bacteria between animals and humans (11).

This study has several limitations that should be considered when interpreting the findings. Participation was voluntary, and recruitment was conducted primarily through online platforms and veterinary clinics, which may have introduced selection bias and may have favoured respondents who are more engaged with veterinary care or more interested in animal health. The study population was predominantly female (79.8%) and included a relatively high proportion of respondents with a medical or veterinary background (24.9%) which together with the convenience-based recruitment strategy, means that the findings may not fully represent the broader pet-owner population in North Macedonia. This may have contributed to the high levels of knowledge and AMR awareness observed in the study, as respondents with a medical/veterinary background demonstrated better knowledge of antibiotic effectiveness and more frequently recognized AMR as a serious health challenge. In addition, the findings were based on self-reported responses and may therefore be influenced by recall bias or social desirability bias, particularly for questions related to antibiotic

use and treatment adherence. Finally, because of the cross-sectional design, the observed associations should not be interpreted as causal relationships.

These findings support the view that AMR is not solely a microbiological phenomenon, but is fundamentally driven by human behaviour (33). Pet-owner knowledge, attitudes, and practices represent critical targets for antimicrobial stewardship efforts within a One Health framework. Addressing behavioural drivers of AMU, including expectations regarding antibiotic therapy, access to antibiotics at home, and treatment-related adherence, may contribute to more responsible antimicrobial use in companion animals. Improved veterinarian–client communication, owner education, and promotion of diagnostic testing may therefore represent important components of antimicrobial stewardship in small animal practice in North Macedonia.

CONCLUSION

In conclusion, this study provides the first data on pet owners' knowledge, attitudes, and practices regarding antimicrobial use and antimicrobial resistance in companion animals in North Macedonia. Despite high general awareness of AMR, important gaps remained in owner understanding and treatment-related behaviour. Reported expectations for antibiotic therapy, use of antibiotics intended for human use in pets, and storage of antibiotics at home suggest that pet owners may play an important role in shaping antimicrobial use in companion animals. These results support the need to include pet owners in antimicrobial stewardship efforts and to strengthen veterinarian–client communication and owner education in small animal practice.

CONFLICT OF INTEREST

The authors declare that they have no financial or non-financial conflict of interest regarding authorship and publication of this article.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the pet owners who participated in this study by completing the questionnaire. We also thank the participating veterinary clinics for their cooperation in facilitating the recruitment of pet owners and the distribution of the questionnaire.

AUTHORS' CONTRIBUTION

IC conceived and designed the study and supervised the manuscript writing. IS, EAP, and MN developed the questionnaire. IS drafted the manuscript and interpreted the results. AC conducted the statistical analysis. BC, JV, MN, MP, and BD contributed to data analysis and interpretation. MN, EAP, and LR participated in manuscript review and editing. All authors reviewed and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

REFERENCES

- World Health Organization (2025). Global antibiotic resistance surveillance report 2025: WHO Global Antimicrobial Resistance and Use Surveillance System (GLASS). Geneva: World Health Organization[A1.1]
<https://doi.org/10.2471/B09585>
- Mader, R., Demay, C., Jouvin-Marche, E., Ploy, M.C., Barraud, O., Bernard, S., et al. (2022). Defining the scope of the European Antimicrobial Resistance Surveillance network in veterinary medicine (EARS-Vet): a bottom-up and One Health approach. *J Antimicrob Chemother.* 77(3): 816-826.
- European Medicines Agency (2025). European sales and use of antimicrobials for veterinary medicine (ESUAvet): annual surveillance report for 2023. Amsterdam: European Medicines Agency[A2.1]
- Zhao, R., Hao, J., Yang, J., Tong, C., Xie, L., Xiao, D., Zeng, Z., Xiong, W. (2022). The co-occurrence of antibiotic resistance genes between dogs and their owners in families. *Imeta.* 1(2): e21.
<https://doi.org/10.1002/imt2.21>
PMid:38868570
- Kovačević, Z., Gračner, G.G., Tomanić, D., Vlahović, K., Šarić, L., Novaković, D., et al. (2025). Compounding and use of human medicinal products in small animal practice: what are the perspectives of veterinarians?-A pilot study. *Vet Sci.* 12(9): 914.
<https://doi.org/10.3390/vetsci12090914>
PMid:41012840 PMCID:PMC12474484
- Marco-Fuertes, A., Marin, C., Lorenzo-Rebenaque, L., Vega, S., Montoro-Dasi, L. (2022). Antimicrobial resistance in companion animals: a new challenge for the One Health approach in the European Union. *Vet Sci.* 9(5): 208.
<https://doi.org/10.3390/vetsci9050208>
PMid:35622736 PMCID:PMC9146952
- European Medicines Agency (2019). Categorisation of antibiotics in the European Union. Amsterdam: European Medicines Agency[A3.1]
- Caneschi, A., Bardhi, A., Barbarossa, A., Zaghini, A. (2023). The use of antibiotics and antimicrobial resistance in veterinary medicine, a complex phenomenon: a narrative review. *Antibiotics (Basel).* 12(3): 487.
<https://doi.org/10.3390/antibiotics12030487>
PMid:36978354 PMCID:PMC10044628
- Hardefeldt, L.Y., Holloway, S., Trott, D.J., Shipstone, M., Barrs, V.R., Malik, R., et al. (2017). Antimicrobial prescribing in dogs and cats in Australia: results of the Australasian infectious disease advisory panel survey. *J Vet Intern Med.* 31(4): 1100-1107.
<https://doi.org/10.1111/jvim.14733>
PMid:28514013 PMCID:PMC5508328
- Shikoska, I., Cvetkovikj, A., Nikolovski, M., Cvetkovikj, I. (2024). Understanding antimicrobial prescription practices: Insights from small animal veterinarians in North Macedonia. *Mac Vet Rev.* 47(2): 103-114.
<https://doi.org/10.2478/macvetrev-2024-0020>
- Monteiro, H.I.G., Silva, V., de Sousa, T., Calouro, R., Saraiva, S., Igrejas, G., Poeta, P. (2025). Antimicrobial resistance in European companion animals practice: a One Health approach. *Animals.* 15(12): 1708.
<https://doi.org/10.3390/ani15121708>
PMid:40564260 PMCID:PMC12189186
- Grönthal, T., Österblad, M., Eklund, M., Jalava, J., Nykäsenoja, S., Pekkanen, K., et al. (2018). Sharing more than friendship - transmission of NDM-5 ST167 and CTX-M-9 ST69 *Escherichia coli* between dogs and humans in a family, Finland, 2015. *Euro Surveill.* 23(27): 1700497.
<https://doi.org/10.2807/1560-7917.ES.2018.23.27.1700497>
PMid:29991384 PMCID:PMC6152158
- Dickson, A., Smith, M., Smith, F., Park, J., King, C., Currie, K., et al. (2019). Understanding the relationship between pet owners and their companion animals as a key context for antimicrobial resistance-related behaviours: an interpretative phenomenological analysis. *Health Psychol Behav Med.* 7(1): 45-61.
<https://doi.org/10.1080/21642850.2019.1577738>
PMid:34040838 PMCID:PMC8114347
- Hopman, N.E.M., Hulscher, M.E.J.L., Graveland, H., Speksnijder, D.C., Wagenaar, J.A., Broens, E.M. (2018). Factors influencing antimicrobial prescribing by Dutch companion animal veterinarians: A qualitative study. *Prev Vet Med.* 158, 106-113.
<https://doi.org/10.1016/j.prevetmed.2018.07.013>
PMid:30220383

15. Smith, M., King, C., Davis, M., Dickson, A., Park, J., Smith, F., et al. (2018). Pet owner and vet interactions: Exploring the drivers of AMR. *Antimicrob Resist Infect Control*. 7, 46.
<https://doi.org/10.1186/s13756-018-0341-1>
PMid:29619213 PMCID:PMC5879597
16. Aithal, S., Guo, H., Teo, B.H., Chua, T., Hildon, Z.J.-L., Chow, A. (2025). Pet owners' knowledge of antibiotic use and antimicrobial resistance and their antibiotic practices: comparison between contexts of self and pet. *Antibiotics (Basel)*. 14(2): 158.
<https://doi.org/10.3390/antibiotics14020158>
PMid:40001402 PMCID:PMC11851957
17. Scarborough, R.O., Sri, A.E., Browning, G.F., Hardefeldt, L.Y., Bailey, K.E. (2023). 'Brave enough': a qualitative study of veterinary decisions to withhold or delay antimicrobial treatment in pets. *Antibiotics (Basel)*. 12(3): 540.
<https://doi.org/10.3390/antibiotics12030540>
PMid:36978407 PMCID:PMC10044613
18. Dias, M.C., Alpizar-Jara, R., Lavrador, C., Marques, C., Broens, E.M., Duarte, E.L. (2024). Companion animal owners' knowledge, attitudes and perceptions regarding antibiotic use in Portugal. *Antibiotics*. 13(6): 533.
<https://doi.org/10.3390/antibiotics13060533>
PMid:38927199 PMCID:PMC11201214
19. Frey, E., Kedrowicz, A., Hedgpeth, M.W. (2022). Exploring companion animal caretakers' attitudes, perceptions and behavioural drivers of antimicrobial use within the social context of veterinary care. *J Small Anim Pract*. 63(12): 873-881.
<https://doi.org/10.1111/jsap.13549>
PMid:36084956 PMCID:PMC10086989
20. Arnecke, A.L., Schwarz, S., Lübke-Becker, A., Jensen, K.C., Bahramsoltani, M. (2024). A Survey on companion animal owners' perception of veterinarians' communication about zoonoses and antimicrobial resistance in Germany. *Animals (Basel)*. 14(22): 3346.
<https://doi.org/10.3390/ani14223346>
PMid:39595398 PMCID:PMC11590884
21. Scarborough, R., Hardefeldt, L., Browning, G., Bailey, K. (2021). Pet owners and antibiotics: knowledge, opinions, expectations, and communication preferences. *Antibiotics (Basel)*. 10(11): 1326.
<https://doi.org/10.3390/antibiotics10111326>
PMid:34827264 PMCID:PMC8615269
22. Redding, L.E., Cole, S.D. (2019). Pet owners' knowledge of and attitudes toward the judicious use of antimicrobials for companion animals. *J Am Vet Med Assoc*. 254(5): 626-635.
<https://doi.org/10.2460/javma.254.5.626>
PMid:30779623
23. Candellone, A., Badino, P., Girolami, F., Ala, U., Mina, F., Odore, R. (2023). Dog owners' attitude toward veterinary antibiotic use and antibiotic resistance with a focus on canine diarrhea management. *Animals (Basel)*. 13(6): 1061.
<https://doi.org/10.3390/ani13061061>
PMid:36978602 PMCID:PMC10044205
24. Stallwood, J., Shirlow, A., Hibbert, A. (2020). A UK-based survey of cat owners' perceptions and experiences of antibiotic usage. *J Feline Med Surg*. 22(2): 69-76.
<https://doi.org/10.1177/1098612X19826353>
PMid:30720397 PMCID:PMC10814557
25. Wang, X., Lin, L., Xuan, Z., Li, L., Zhou, X. (2018). Keeping antibiotics at home promotes self-medication with antibiotics among Chinese university students. *Int J Environ Res Public Health*. 15(4): 687.
<https://doi.org/10.3390/ijerph15040687>
PMid:29621165 PMCID:PMC5923729
26. Allerton, F., Prior, C., Bağcıgil, A.F., Broens, E.M., Callens, B., Damborg, P., et al. (2021). Overview and evaluation of existing guidelines for rational antimicrobial use in small-animal veterinary practice in Europe. *Antibiotics (Basel)*. 10(4): 409.
<https://doi.org/10.3390/antibiotics10040409>
PMid:33918617 PMCID:PMC8069046
27. Galarce, N., Dumont-Viollaz, A., Longa, J., del Río, L.C., Núñez, A., Guzmán-Marín, B., Thomson, P. (2025). Perceptions of antimicrobial use and resistance among pet owners in Chile: A cross-sectional One Health survey. *Vet World*. 18(8): 2450-2459.
<https://doi.org/10.14202/vetworld.2025.2450-2459>
PMid:41064845 PMCID:PMC12501537
28. Wright, E., Jessen, L.R., Tompson, A., Rutland, C., Singleton, D., Battersby, I., et al. (2024). Influencing attitudes towards antimicrobial use and resistance in companion animals-the impact on pet owners of a short animation in a randomized controlled trial. *JAC Antimicrob Resist*. 6 (3): dlac065.
<https://doi.org/10.1093/jacamr/dlae065>
PMid:38716404 PMCID:PMC11073752
29. Tomanic, D., Stojanovic, D., Belić, B., Davidov, I., Novakov, N., Radinovic, M., et al. (2021). Off label use of human approved drugs in treatment of dogs in the Republic of Serbia. *Pol J Vet Sci*. 24(3): 399-407.
<https://doi.org/10.24425/pjvs.2021.138731>
PMid:34730304

30. Shikoska, I., Duvnjak, S., Koritnik, T., Chapkunovska, B., Vlahov, J., Ratkova Manovska, M., Cvetkovikj, A., Cvetkovikj, I. (2025). Extended-spectrum β -Lactamase-/AmpC-Producing *Escherichia coli* and associated risk factors in shelter dogs: a baseline study in North Macedonia. *Microbiol Res.* 16(9): 206. <https://doi.org/10.3390/microbiolres16090206>
31. Shikoska, I., Hristovska, Z.P., Matevski, I., Pavlova, M.J., Manovska, M.R., Cvetkovikj, A., Cvetkovikj, I. (2025). Phenotypic and molecular characterization of antimicrobial resistance in canine *Staphylococci* from North Macedonia. *Mac Vet Rev.* 48 (2): 159-172. <https://doi.org/10.2478/macvetrev-2025-0022>
32. Stein, M.R., Weese, J.S., Stull, J.W., McClure, J.T., Evason, M. (2022). Knowledge, attitudes and influencers of cat owners in North America around antimicrobials and antimicrobial stewardship. *J Feline Med Surg.* 24(6): e90-e97. <https://doi.org/10.1177/1098612X221090456>
33. Chadborn, T., Williams, S., Jorgensen, T., Price, C., Buckel, A., Altieri, E. (2023). An approach for embedding behavioural science in antimicrobial resistance One Health research. *J Infect Public Health.* 16 (Suppl 1): 134-140. <https://doi.org/10.1016/j.jiph.2023.11.001> PMID:37973498

Please cite this article in press as: Shikoska I., Vlahov J., Dimitrievski B., Chapkunovska B., Rashikj Lj., Prodanov M., Nikolovski M., Atanaskova Petrov E., Cvetkovikj A., Cvetkovikj I. Pet owners' knowledge, attitudes, and practices regarding antimicrobial use and antimicrobial resistance in companion animals in North Macedonia. *Mac Vet Rev* 2026; 49 (2): i-xiv. <https://doi.org/10.2478/macvetrev-2026-0026>