



## MICROBIOLOGICAL QUALITY OF SOFT, SEMI-HARD AND HARD CHEESES DURING THE SHELF-LIFE

Josip Vrdoljak<sup>1</sup>, Vesna Dobranić<sup>2</sup>, Ivana Filipović<sup>2</sup>, Nevijo Zdolec<sup>2</sup>

<sup>1</sup>PUDA D.O.O., Economic Zone Čaporice 133, 21240 Čaporice, Croatia

<sup>2</sup>Department of Hygiene, Technology and Food Safety,  
Faculty of Veterinary Medicine, University of Zagreb,  
Heinzelova 55, 10000 Zagreb, Croatia

Received 1 July 2015; Received in revised form 3 November 2015; Accepted 19 November 2015

### ABSTRACT

Cheeses as ready-to-eat food should be considered as a potential source of foodborne pathogens, primarily *Listeria monocytogenes*. The aim of present study was to determine the microbiological quality of soft, semi-hard and hard cheeses during the shelf-life, with particular reference to *L. monocytogenes*. Five types of cheeses were sampled at different time-points during the cold storage and analyzed for presence of *Salmonella* and *L. monocytogenes*, as well as lactic acid bacteria, *Escherichia coli*, coagulase-positive staphylococci, yeasts, molds, sulfite-reducing clostridia and *L. monocytogenes* counts. Water activity, pH and NaCl content were monitored in order to evaluate the possibility of *L. monocytogenes* growth. Challenge test for *L. monocytogenes* was performed in soft whey cheese, to determine the growth potential of pathogen during the shelf-life of product. All analyzed cheeses were compliant with microbiological criteria during the shelf-life. In soft cheeses, lactic acid bacteria increased in the course of the shelf-life period (1.2-2.6 log increase), while in semi-hard and hard cheeses it decreased (1.6 and 5.2 log decrease, respectively). Soft cheeses support the growth of *L. monocytogenes* according to determined pH values (5.8-6.5), water activity (0.99-0.94), and NaCl content (0.3-1.2%). Challenge test showed that *L. monocytogenes* growth potential in selected soft cheese was 0.43 log<sub>10</sub> cfu/g during 8 days at 4°C. Water activity in semi-hard and hard cheeses was a limiting factor for *Listeria* growth during the shelf-life. Soft, semi-hard and hard cheeses were microbiologically stable during their defined shelf-life. Good manufacturing and hygienic practices must be strictly followed in the production of soft cheeses as *Listeria*-supporting food and be focused on preventing (re)contamination.

**Key words:** cheese, shelf-life, *L. monocytogenes*

### INTRODUCTION

Microbiological stability and safety of food during storage is related to many factors. Ready-to-eat food products, including cheeses, are intended for consumption without any treatment between final production step and consumption. The course of microbiological changes in different cheeses during storage and shelf-life depends on the production technology and cheese type (pasteurization, starters, acidity, ripening, etc.), physico-chemical properties

(acidity, moisture content, salt content), storage conditions (microclimate, handling, packaging). *Listeria monocytogenes* is the most important foodborne pathogen in cheese microbiology, mainly in the post-processing phase (5). European Regulation on microbiological criteria of food (EC 2073/2005) categorizes foods into two groups, supporting or not supporting the growth of *L. monocytogenes*. Foods with pH ≤4.4 or water activity ( $a_w$ ) ≤0.92, or products with pH ≤5.0 and  $a_w$  ≤0.94, as well as products with shelf-life less than 5 days are considered as safe regarding *Listeria* outgrowth (4). Recent survey conducted by EFSA showed that 0.06% of soft and semi-soft cheeses (n=3452) are exceeding the level of 100 cfu/g at the end of shelf-life. The occurrence of *L. monocytogenes* in cheese samples was 0.47% (6). In Croatia, microbiological surveys in recent years were mainly performed on traditional dairy products from non-pasteurized milk, and *L. monocytogenes* was frequently present (7, 12). Thus, the aim of the

Corresponding author: Assist. Prof. Nevijo Zdolec, PhD

E-mail address: [nzdolec@vef.hr](mailto:nzdolec@vef.hr)

Present address: Department of Hygiene, Technology and Food Safety,  
Faculty of Veterinary Medicine, University of Zagreb,  
Heinzelova 55, 10000 Zagreb, Croatia  
Phone: ++38512390192

**Copyright:** © 2015 Vrdoljak J. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Competing Interests:** The authors have declared that no competing interests exist.

**Available Online First:** 4 December 2015

**Published on:** 15 March 2016

<http://dx.doi.org/10.1515/macvetrev-2015-0068>

present study was to evaluate the microbiological quality of pasteurized-milk cheeses during the shelf-life with particular reference to *L. monocytogenes* prevalence and counts. The growth potential of *L. monocytogenes* was also evaluated in soft whey cheese during the shelf-life.

## MATERIAL AND METHODS

Soft, semi-hard and hard cheeses were produced according to standard procedure in a local cheese producing plant and sampled at the end of production. Ten units of each cheese were sampled, stored in laboratory at 4°C and periodically analyzed during defined shelf-life (Table 1).

pH with pH-meter (pH 510, Eutech instruments, Netherlands) and NaCl following Mohr's method. Challenge test was performed in whey cheese "skuta" (shelf-life 8 days) inoculated with mixed culture of *L. monocytogenes* (two strains of cheese origin and *L. monocytogenes* ATCC 7644). The strains were grown in Brain Heart Infusion Broth (bioMerieux, France) at 37°C for 24 h. One ml of each culture was centrifuged at 10000 g for 10 min, followed by supernatant removal and washing the cells with sterile saline water. Cells were diluted in 1 ml of sterile saline water and serially diluted to determine the cell number, using PALCAM agar (Merck, Germany) incubated at 37°C for 24-48 h.

**Table 1.** Sampling scheme and parameters monitored in cheese during shelf-life

Cheese type	Cheese name	Shelf-life	Time-point sampling	Microbiological parameters	Chemical parameters
Soft cheese	Whey cheese Skuta	8 days	Day 5, 7, 8	<i>Salmonella</i> , <i>L.monocytogenes</i> , <i>E.coli</i> , <i>S. aureus</i> ,	Water activity, pH, NaCl
	Škripavac	15 days	Day 6, 9, 12, 15	yeasts, molds, lactic acid bacteria	
Semi-hard cheese	Trappist	9 months	Month 3,4,6,9	<i>Salmonella</i> , <i>L.monocytogenes</i> , <i>E.coli</i> , <i>S. aureus</i> , Sulfite- reducing clostridia, lactic acid bacteria	
Hard cheese	Livanjski	15 months	Month 3,4,6,11,12,15	<i>Salmonella</i> , <i>L.monocytogenes</i> , <i>E.coli</i> , <i>S. aureus</i> ,	
	Hard cheese in olive oil	9 months	Month 3,5,8,9	Sulfite- reducing clostridia, lactic acid bacteria	

For microbiological analyses, 25 g of sample was diluted in 225 ml of appropriate media (Buffered Peptone Water, Half-Fraser Broth or Peptone salt water) and homogenized for 2 min (Stomacher, Sedward, UK). Serial decimal dilutions were prepared and 0.1 ml or 1 ml of selected dilution were used for evaluation of lactic acid bacteria count (MRS, Merck, Darmstadt, Germany) at 30°C for 48 h, *Staphylococcus aureus* (Baird Parker agar, Merck, Germany) at 37°C for 48 h, *Escherichia coli* (Rapid E. coli, Bio-Rad, France) 37/44°C for 24 h, and sulfite-reducing clostridia (Iron Sulphyte Agar, bioMerieux, Marcy l'Etoile, France). *Salmonella* spp. was determined following a standard ISO 6579 method and Molecular Detection System (MDS, 3M, USA). *L. monocytogenes* presence and count was determined using ISO 11290-1 and 11290-2, respectively.

Water activity ( $a_w$ ) was determined by means of HygroPalm AW1 (Rotronic, Switzerland),

Appropriate dilutions were taken for cheese portion inoculation to obtain an inoculation level of 30-50 cfu/g. Cheese samples were inoculated in triplicate, vacuum-packed and stored at 4°C for 8 days. *L. monocytogenes* count, pH and water activity were determined at day 0 and at the end of shelf-life.

## RESULTS

The results of the microbiological and chemical analyses of cheeses during the storage are summarized in Table 2. Foodborne pathogens *Salmonella* spp. and *L. monocytogenes* were absent in all cheese samples analyzed during the shelf-life. *Escherichia coli*, *L. monocytogenes*, sulfite-reducing clostridia and *S. aureus* were below the detection limits of the methods used at all sampling points.

*L. monocytogenes* counts were below 10 cfu/g in all cheese samples during storage. Population of lactic acid bacteria increased during the storage of soft cheeses, reaching 7-8 log cfu/g at the end of shelf-life. The pH values, water activity and NaCl content in soft cheeses were in the range of *Listeria*-supporting values. Semi-hard and hard cheeses were characterized by decrease of lactic acid bacteria counts during the storage and a slight pH increase. The pH values and NaCl content found in semi-hard and hard cheeses were not a limiting factor for *Listeria* growth. However, it is evident that water activity was below values that enables the growth of most foodborne pathogens.

Since soft cheeses showed to be supporting of *Listeria* growth according to their physico-chemical

characteristics, the challenge test was performed with *L. monocytogenes*. Results of *L. monocytogenes* growth potential in whey cheese skuta are shown in Table 3, as well as pH and  $a_w$  values (Table 4). Despite high water activity and optimal pH, *L. monocytogenes* inoculated in low numbers (30-50 cfu/g) didn't reach a critical limit of 100 cfu/g in cheese stored at 4 °C for 8 days.

## DISCUSSION

Microbiological quality and interpretation of microbiological findings of different kind of cheeses depends on the sampling points during the production or retail phase. In present study, the

**Table 2.** Results of microbiological and physico-chemical analyses of cheeses during the shelf-life

Cheese type	Cheese name	Microbiological parameter and sampling day	Microbial counts (log <sub>10</sub> cfu/g)	Results of physico-chemical analyses		
				pH	$a_w$	NaCl (%)
Soft cheese	Whey cheese Skuta	Lactic acid bacteria				
		Day 5	4.20±0.07	6.21±0.02	0.945±0.01	0.12±0.02
		Day 7	4.60±0.05	6.53±0.01	0.960±0.01	0.23±0.03
		Day 8	6.80±0.02	5.80±0.02	0.990±0.01	0.29±0.04
		Yeasts				
		Day 5	2.47±0.01			
	Škripavac	Day 7	2.77±0.11			
		Day 8	2.95±0.12			
		Lactic acid bacteria				
		Day 6	6.62±0.15	6.28±0.02	0.950±0.02	1.10±0.02
Semi-hard cheese	Trappist	Day 9	6.50±0.07	6.40±0.03	0.960±0.01	1.10±0.02
		Day 12	7.25±0.12	6.55±0.03	0.978±0.02	1.20±0.01
		Day 15	7.82±0.14	6.20±0.01	0.990±0.02	1.22±0.02
		Yeasts				
		Day 6	2.30±0.11			
		Day 9	2.54±0.06			
	Livnjski	Day 12	2.58±0.07			
		Day 15	2.77±0.10			
		Lactic acid bacteria				
		Day 90	6.20±0.12	5.40±0.04	0.910±0.01	1.51±0.02
Hard cheese	Trappist	Day 120	5.04±0.10	5.48±0.02	0.900±0.04	1.68±0.03
		Day 180	4.80±0.08	5.80±0.04	0.866±0.03	2.12±0.06
		Day 270	4.62±0.10	5.96±0.02	0.806±0.02	2.85±0.03
	Livnjski	Lactic acid bacteria				
		Day 90	7.20±0.15	5.52±0.03	0.900±0.02	1.85±0.02
		Day 120	5.83±0.10	5.60±0.02	0.907±0.05	2.20±0.02
		Day 180	4.60±0.12	5.72±0.05	0.880±0.03	2.28±0.04
		Day 330	3.40±0.12	5.80±0.02	0.795±0.03	3.30±0.01
		Day 360	2.20±0.05	5.92±0.02	0.791±0.06	3.35±0.02
	Hard cheese in olive oil	Day 450	2.00±0.02	5.97±0.02	0.650±0.04	3.62±0.03
		Lactic acid bacteria				
		Day 90	6.80±0.11	5.72±0.02	0.890±0.02	1.74±0.02
		Day 150	6.78±0.14	5.80±0.04	0.870±0.03	1.75±0.02
	Hard cheese in olive oil	Day 240	6.75±0.05	5.84±0.03	0.842±0.04	1.80±0.04
		Day 270	6.42±0.04	5.90±0.03	0.760±0.04	1.92±0.03

**Table 3.** Growth potential of *Listeria monocytogenes* in whey cheese skuta

Batch	Sample	<i>L. monocytogenes</i> number (log <sub>10</sub> cfu/g) Day 0	<i>L. monocytogenes</i> number (log <sub>10</sub> cfu/g) Day 8	Growth potential (log <sub>10</sub> cfu/g)
<b>I</b>	1a	1	1.84	0.24
	1b	1.60	1.90	
	1c	1.69	1.69	
<b>II</b>	2a	1.47	1.90	0.43
	2b	1.30	1.95	
	2c	1.69	1.95	
<b>III</b>	3a	1.30	1.69	0.09
	3b	1.60	1.30	
	3c	1.69	1.90	

**Table 4.** pH and aw in whey cheese skuta during challenge test

Batch	Sample	Day 0		Day 8	
		pH	a <sub>w</sub>	pH	a <sub>w</sub>
<b>I</b>	1a	6.22	0.952	5.80	0.992
	1b	6.30	0.950	5.48	0.996
	1c	6.06	0.955	5.62	0.998
<b>II</b>	2a	6.42	0.960	5.45	0.993
	2b	6.23	0.958	5.50	0.997
	2c	6.30	0.950	5.62	0.998
<b>III</b>	3a	6.17	0.952	5.50	0.998
	3b	6.15	0.980	5.46	0.997
	3c	6.22	0.976	5.52	0.994

phase of post-processing was monitored in order to evaluate compliance with microbiological criteria, with particular reference to *L. monocytogenes* counts. In relation to overall microbiological quality, results showed that soft, semi-hard and hard cheeses were microbiologically stable during their defined shelf-life. Microbiological stability of semi-hard and hard cheeses relies on several hurdles including starter cultures competitiveness, increased salt content and, most importantly, low water activity (2, 9). Physico-chemical values and their changes during the storage of semi-hard and hard cheeses in our study are in line with previous reports (11, 13, 15). *L. monocytogenes* was not present in viable counts, however it should be stressed that secondary contamination is possible during the cutting of the cheese into quarters or halves, followed by packaging. Prevention of contamination with *Listeria* at this point should be based on Good Hygienic Practice and verification of sanitation programs.

Soft cheeses in general are of limited durability, because of high moisture content and high pH which support the proliferation of some spoilage microorganism like *Enterobacter* spp.

or *Pseudomonas* spp. under cold storage (8). Whey cheese (skuta) used in our study showed physicochemical characteristics comparable with other similar products from the Mediterranean area (1, 10, 16). During the storage of whey cheese skuta the pH decreased and lactic acid bacteria count increased which is in accordance with other studies (10, 16). Lactic acid bacteria are known to be beneficial and functional microbes in many different dairy products (20), however their outgrowth in this kind of products (whey cheese) could contribute to spoilage (8). Soft cheeses are ready-to-eat products that support the growth of *L. monocytogenes* based on their physico-chemical characteristics (14, 18), which is also presented by the results in current survey (pH values 5.8-6.5, water activity 0.99-0.94 and NaCl content 0.3-1.2%). Challenge test showed that *L. monocytogenes* growth potential in whey cheese was 0.43 log<sub>10</sub> cfu/g during 8 days at 4°C, meaning that the proposed shelf-life is acceptable for products stored under defined conditions. Many studies from last decades emphasized that *L. monocytogenes* represents a serious public-health problem due to high prevalence in soft cheeses (7, 17, 19). Recent studies are more focused on

bacterial kinetics in cheeses made from pasteurized milk during their shelf-life at different storage conditions, and related application of bio-protective strategies (3).

## CONCLUSION

Compliance with microbiological criteria at the end of cheese production (final product) doesn't guarantee that microbiological hazards are excluded. *Listeria monocytogenes* is a ubiquitous bacteria and secondary contamination of products is possible under poor hygienic conditions. Despite the fact that the growth of the pathogen is limited in semi-hard and hard cheeses by low water activity, the secondary (surface) contamination could result in hazardous products. The significance of following strict hygienic procedures is evident even more in soft cheese production, since they support the growth of *L. monocytogenes*.

## REFERENCES

1. Antunac, N., Hudik, S., Mikulec, N., Maletić, M., Horvat, I., Radeljević, B., Havranek, J. (2011). Proizvodnja i kemijski sastav Istarske i Paške skute. *Mljekarstvo* 61, 326-335. (in Croatian)
2. Bishop, J.R., Smukowski, M. (2006). Storage temperatures necessary to maintain cheese safety. *Food Protect. Trends* 26, 714-724.
3. Coelho, M.C., Silva, C.C.G., Ribeiro, S.C., Dapkevicius, M.L.N.E., Rosa, H.J.D. (2014). Control of *Listeria monocytogenes* in fresh cheese using protective lactic acid bacteria. *Int. J. Food Microbiol.* 191, 53-59.  
<http://dx.doi.org/10.1016/j.ijfoodmicro.2014.08.029>  
PMid:25222327
4. Commission regulation (EC) No 2073/2005 on microbiological criteria for foods. Official Journal of the European Union L338/1-338/26.
5. Donnelly, C.W. (2004). Growth and survival of microbial pathogens in cheese. In: Fox, P.F., McSweeney, P.L.H., Cogan, T.M., Guinee, T. P. (Eds.), *Cheese - Chemistry, Physics and Microbiology*, Third Edition – Vol. 1: general aspects. (pp 541-559), London: Elsevier Ltd.  
[http://dx.doi.org/10.1016/s1874-558x\(04\)80081-2](http://dx.doi.org/10.1016/s1874-558x(04)80081-2)
6. EFSA (2013). Analysis of the baseline survey on the prevalence of *Listeria monocytogenes* in certain ready-to-eat foods in the EU, 2010-2011 Part A: *Listeria monocytogenes* prevalence estimates. *EFSA Journal* 11, 3241.
7. Kozačinski, L., Hadžiosmanović, M. (2001). The occurrence of *Listeria monocytogenes* in home-made dairy products. *Tierartz Umsch.* 56, 590-594.
8. Ledenbach, L.H., Marshall, L.T. (2009). Microbiological spoilage of dairy products. In: Sperber, W.H., Doyle, M.P. (Eds.), *Compendium of the microbiological spoilage of foods and beverages* (pp 41-67), Springer Science.  
[http://dx.doi.org/10.1007/978-1-4419-0826-1\\_2](http://dx.doi.org/10.1007/978-1-4419-0826-1_2)
9. Leong, W.M., Geier, R., Engstrom, S., Ingham, S., Ingham, B., Smukowski, M. (2014). Growth of *Listeria monocytogenes*, *Salmonella* spp., *Escherichia coli* O157:H7 and *Staphylococcus aureus* on cheese during extended storage at 25 °C. *J. Food Protect.* 77, 1275-1288.  
<http://dx.doi.org/10.4315/0362-028X.JFP-14-047>  
PMid:25198588
10. Litopoulou-Tzanetaki, E., Tzanetakis, N. (2011). Microbiological characteristics of Greek traditional cheese. *Small Rum Res.* 101, 17-32.  
<http://dx.doi.org/10.1016/j.smallrumres.2011.09.022>
11. Marijan, A., Džaja, P., Bogdanović, T., Škoko, I., Cvetnić, Ž., Dobranić, V., Zdolec, N., Šatrović, E., Sverin, K. (2014). Influence of ripening time on the amount of certain biogenic amines in rind and core of cow milk Livno cheese. *Mljekarstvo* 64, 159-169.  
<http://dx.doi.org/10.15567/mljekarstvo.2014.0303>
12. Markov, K., Frece, J., Čvek, D., Delaš, F. (2009). *Listeria monocytogenes* and other contaminants in fresh cheese and cream from Zagreb city area domestic production. *Mljekarstvo* 59, 225-231.
13. Matić, A., Kalit, S., Salajpal, K., Ivanković, S., Sarić, Z. (2014). Consumers' preferences and composition of Livanjski cheese in relation to its sensory characteristics. *Mljekarstvo* 64, 170-177.  
<http://dx.doi.org/10.15567/mljekarstvo.2014.0304>
14. Melo, J., Andrew, P.W., Faleiro, M.L. (2015). *Listeria monocytogenes* in cheese and the dairy environment remains a food safety challenge: The role of stress responses. *Food Res. Int.* 67, 75-90.  
<http://dx.doi.org/10.1016/j.foodres.2014.10.031>
15. Merćep, A., Kirin, S., Zdolec, N., Cvrtila Fleck, Ž., Filipović, I., Njari, B., Mitak, M., Kozačinski, L. (2010). Quality of Trappist cheese from Croatian dairy plant. *Mljekarstvo* 60, 288-298.
16. Papaioannou, G., Chouliara, I., Karatapanis, A.E., Kontominas, M.G., Savvaidis, I.N. (2007). Shelf-life of a Greek whey cheese under modified atmosphere packaging. *Int. Dairy J.* 17, 358-364.  
<http://dx.doi.org/10.1016/j.idairyj.2006.04.001>
17. Pintado, C.M.B.S., Oliveira, A., Pampulha, M.E., Ferreira, M.A.S.S. (2005). Prevalence and characterization of *Listeria monocytogenes* isolated from soft cheese. *Food Microbiol.* 22, 79-85.  
<http://dx.doi.org/10.1016/j.fm.2004.04.004>

18. Rosshaug, P.S., Detmer, A., Ingmer, H., Larsen, M.H. (2012). Modeling the growth of *Listeria monocytogenes* in soft Blue-White cheeses. *Appl. Environ. Microbiol.* 78, 8508-8514.  
<http://dx.doi.org/10.1128/AEM.01865-12>  
PMid:22983971 PMCID:PMC3502931
19. Rudolf, M., Scherer, S. (2001). High incidence of *Listeria monocytogenes* in European red smear cheese. *Int. J. Food Microbiol.* 63, 91-98.  
[http://dx.doi.org/10.1016/S0168-1605\(00\)00413-X](http://dx.doi.org/10.1016/S0168-1605(00)00413-X)
20. Zdolec, N., Lazić, S., Kozačinski, L., Hadžiosmanović, M., Filipović, I. (2007). The inhibitory activity of lactic acid bacteria isolated from fresh cow cheese. *Mljekarstvo* 57, 5-13.